

A neurophysiological deficit in early visual processing in schizophrenia patients with auditory hallucinations

JÜRGEN KAYSER,^{a,b} CRAIG E. TENKE,^{a,b} CHRISTOPHER J. KROPPMANN,^b DANIEL M. ALSCHULER,^b SHIVA FEKRI,^b ROBERTO GIL,^{a,c} L. FREDRIK JARSKOG,^{a,d} JILL M. HARKAVY-FRIEDMAN,^a AND GERARD E. BRUDER^{a,b}

^aDepartment of Psychiatry, Columbia University College of Physicians and Surgeons, New York, New York, USA

^bDivision of Cognitive Neuroscience, New York State Psychiatric Institute, New York, New York, USA

^cClinical Psychobiology, New York State Psychiatric Institute, New York, New York, USA

^dDepartment of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA

Abstract

Existing 67-channel event-related potentials, obtained during recognition and working memory paradigms with words or faces, were used to examine early visual processing in schizophrenia patients prone to auditory hallucinations (AH, $n = 26$) or not (NH, $n = 49$) and healthy controls (HC, $n = 46$). Current source density (CSD) transforms revealed distinct, strongly left- (words) or right-lateralized (faces; N170) inferior-temporal N1 sinks (150 ms) in each group. N1 was quantified by temporal PCA of peak-adjusted CSDs. For words and faces in both paradigms, N1 was substantially reduced in AH compared with NH and HC, who did not differ from each other. The difference in N1 between AH and NH was not due to overall symptom severity or performance accuracy, with both groups showing comparable memory deficits. Our findings extend prior reports of reduced auditory N1 in AH, suggesting a broader early perceptual integration deficit that is not limited to the auditory modality.

Descriptors: Schizophrenia, Auditory hallucination, Event-related potentials (ERPs), Visual N1, Current source density (CSD), Principal components analysis (PCA)

Impairments of working memory (WM) and recognition memory (RM) are core features in patients with schizophrenia (Barch, 2005). It has been suggested that impairments in early sensory, bottom-up processing contribute to the WM deficit in schizophrenia (Javitt, 2009). Javitt, Shelley, Silipo, and Lieberman (2000) found evidence that schizophrenia is associated with impairments of auditory sensory (echoic) memory, which is evident even in simple tone discrimination tasks. A deficit in precision of encoding physical stimulus properties can, in turn, produce downstream deficits in cognitive tasks involving utilization of memory traces. Studies have successfully exploited the fine temporal resolution of event-related brain potentials (ERPs) to provide direct measures of the time course of ongoing brain activity during memory processing (e.g., Kayser et al., 2006), which may provide further insight into the stage of processing that underlies memory deficits in schizophrenia. ERP components occurring at different latencies

reflect the spatiotemporal sequence of cortical processing, ranging from early sensory perception and attention (e.g., P1 or N1) to later cognitive evaluation (e.g., P3). For instance, Haenschel et al. (2007) found that a deficit in early visual processing, as indexed by a reduction of the P1 component peaking about 100 ms after stimulus onset, contributed to the impaired visual WM performance.

Reductions of early visual (Bruder et al., 1998; Butler et al., 2007; Doniger, Foxe, Murray, Higgins, & Javitt, 2002; Foxe, Doniger, & Javitt, 2001; Neuhaus et al., 2011; Yeap et al., 2006) and auditory (Foxe et al., 2011; Kayser et al., 2001; Rosburg, Boutros, & Ford, 2008; Salisbury, Collins, & McCarley, 2010) ERP components in schizophrenia are often considered as evidence of sensory, bottom-up processing deficits, which may represent an endophenotype for schizophrenia (Foxe et al., 2001, 2011). This has primarily been found in simple target detection tasks, which involve minimal cognitive demand. In fact, task or stimulus processing requirements are of critical importance and may account for conflicting findings concerning early visual processing deficits in schizophrenia (Butler et al., 2007; Foxe et al., 2001). For example, Doniger et al. observed reduced P1 during the recognition of fragmented images while N1 was preserved, whereas Bruder et al. (1998) found reduced N1 but no P1 reduction during a visuospatial discrimination task. Less is known as to whether electrophysiological indices of early sensory processing are

This research was supported by Grant MH066597 from the National Institute of Mental Health (NIMH). We thank Charles L. Brown, III, for developing, providing, and improving excellent waveform plotting software. Preliminary analyses of these data were presented at the 50th Annual Meeting of the *Society for Psychophysiological Research (SPR)*, Portland, Oregon, September 29–October 3, 2010.

Address correspondence to: Jürgen Kayser, New York State Psychiatric Institute, Division of Cognitive Neuroscience, Unit 50, 1051 Riverside Drive, New York, NY 10032, USA. E-mail: kayserj@nyspi.columbia.edu

reduced in schizophrenia during more demanding memory tasks. In three ERP studies (Kayser et al., 1999, 2010; Kayser, Tenke, Gil, & Bruder, 2009), in which schizophrenia patients showed poorer RM for words compared to healthy controls, N1 amplitude tended to be smaller in patients, but this group difference did not attain statistical significance.

There is evidence that N1 amplitude may vary with the severity of patients' symptoms (Rosburg et al., 2008). In particular, reduction of the N1 component to tones has been reported for psychotic patients during periods of auditory hallucinations (Hubl, Koenig, Strik, Garcia, & Dierks, 2007), which could stem from central acoustic interference associated with internal speech (Ford et al., 2001), that is, auditory hallucinations may interfere with processing of tones or spoken words. This raises the possibility that patients having auditory hallucinations may be particularly likely to show reduced N1 amplitude and that an impairment in early auditory processing could contribute to a heightened WM deficit. In a study measuring auditory WM in a word serial position test, which involves storage and rehearsal of phonological and sequential information, patients who reported experiencing auditory hallucinations showed significantly poorer performance than patients who did not report having hallucinations (Bruder et al., 2011). In a multisite neuroimaging study, patients with auditory hallucinations, when compared to nonhallucinating patients, showed decreased activity in left superior temporal and inferior parietal regions to probe stimuli in a Sternberg WM task (Wible et al., 2009). Although stimuli were visually presented numbers, the auditory verbal language system would be expected to be involved in item rehearsal.

Several lines of evidence suggest that auditory verbal hallucinations involve the left temporal lobe and reduced language lateralization (see Badcock & Hugdahl, 2012, for a review). This hypothesis has received support from functional magnetic resonance imaging studies showing reduced activation in left superior-temporal and inferior-parietal regions in hallucinators (Ford et al., 2009; Hugdahl et al., 2007; Wible et al., 2009) and dichotic listening studies, which have consistently found hallucinations to be associated with reduced left hemisphere advantage for words or syllables (Bruder et al., 1995; Green, Hugdahl, & Mitchell, 1994; Hugdahl et al., 2008). Moreover, several electrophysiologic studies using rather heterogenous approaches to study auditory hallucinations have also implicated the left superior temporal cortex (see van Lutterveld, Sommer, & Ford, 2011, for a review). For example, using LORETA as an inverse source localization technique (e.g., Pascual-Marqui, Michel, & Lehmann, 1994), Hubl et al. (2007) reported that the N1 reduction to tones during periods of auditory hallucinations was primarily associated with reduced current density sources in the left temporal lobe. It would therefore be of interest to more directly study lateralized neurophysiologic processes in schizophrenia by employing stimuli known to differentially engage the left or right hemisphere in early perceptual processing (cf. Kayser et al., 2001).

This report takes advantage of ERP data recorded from a large sample of schizophrenia patients and healthy controls during working memory and recognition memory tasks using words and faces (Kayser et al., 2010) to examine whether patients' propensity to experience auditory hallucinations affects early visual processing (cf. van Lutterveld et al., 2011). Given the above findings, we predicted that patients who report having auditory hallucinations would show reduced N1 when compared to patients without hallucinations and healthy controls. If patients who are prone to having auditory hallucinations show reduced N1 amplitude during these visual RM and WM paradigms, this would indicate that it is

not merely due to direct acoustical interference (i.e., hearing voices), and not limited to the auditory modality, thereby suggesting a broader, and modality-unspecific, cognitive deficit. Alternatively, we may observe reduced early visual ERP components in schizophrenia independent of the tendency to experience auditory hallucinations, which would provide further evidence that disrupted cognition in schizophrenia is caused by a fundamental deficit in bottom-up processes (e.g., Javitt, 2009).

Because ERPs depend on the electroencephalogram (EEG) reference location and the operational definition of an ERP component measure, current source density (CSD) and temporal principal components analysis (PCA) were combined to obtain a reference-free estimate of visual N1 (Kayser & Tenke, 2006a), which is strongly left-lateralized for words (Kayser, Tenke, Gates, & Bruder, 2007; Kayser et al., 2009, 2010). Given the involvement of left temporoparietal cortex in word processing (Price, 2000, 2010) and auditory hallucinations (e.g., Badcock & Hugdahl, 2012; van Lutterveld et al., 2011; Wible et al., 2009), we also predicted that the N1 reduction in hallucinators would be most evident over left temporoparietal sites for words.

Methods

Participants

We analyzed the ERP data of 58 inpatients and 17 outpatients (42 male) at New York State Psychiatric Institute and 46 healthy adults (20 male) who had participated in at least one of two visual memory paradigms. Most participants performed both RM (Session 1) and WM (Session 2) tasks at two different recording sessions within a week, typically 1–2 days apart. Left-handed individuals (Oldfield, 1971) and those with a history of neurological illness or substance abuse, as well as participants performing at or below chance level, were excluded. Most importantly, only patients for whom clinical ratings of auditory hallucinations were available for the time of ERP testing were included. The auditory hallucination item of the Scale for the Assessment of Positive Symptoms (SAPS; Andreasen, 1984) was used to form symptom subgroups by separating patients who reported experiencing auditory hallucinations (AH) in the past week (SAPS auditory hallucination item ≥ 1) from those who did not (NH).

Table 1 details the demographic and clinical variables, DSM-IV (American Psychiatric Association, 1994) criteria, and medication status for the two patient subgroups and healthy controls (HC) in the two memory paradigms. Most patients (14 AH, 30 NH) and 42 healthy adults provided data for both memory paradigms. Although the groups did not differ in gender, age (range 18–56 years) or degree of right-handedness (laterality quotient range 20–100), patients had significantly less education than controls. However, the available verbal IQ data (WAIS) suggested that the patients' verbal skills were well within normal range and did not differ between AH and NH subgroups. Although hallucinators and non-hallucinators did not differ in age of onset (range 9–37 years) or illness duration (range 0–47 years), the total score of the Brief Psychiatric Rating Scale (BPRS), which was derived from the Positive and Negative Syndrome Scale (PANSS; Kay, Opler, & Fishbein, 1992), indicated that hallucinators were more disturbed than nonhallucinators. This difference is also reflected in higher positive and negative PANSS subscales in hallucinators than non-hallucinators. Notwithstanding, all electrophysiologic findings reported were not affected by subgroup differences in total BPRS measures of severity (see below).

Table 1. Means (SD) for Demographic and Clinical Variables, DSM-IV Criteria, Medication Status, and Sample Sizes for Each Paradigm

Paradigm	Recognition memory (RM)			Working memory (WM)		
	AH (<i>n</i> = 19)	NH (<i>n</i> = 32)	HC (<i>n</i> = 44)	AH (<i>n</i> = 21)	NH (<i>n</i> = 47)	HC (<i>n</i> = 44)
Group ^a						
Gender (male/female)	10/9	18/14	18/26	12/9	26/21	19/25
Age (years)	29.8 (10.2)	28.6 (8.0)	26.2 (6.2)	30.6 (11.0)	28.8 (8.0)	26.4 (6.6)
Education (years) ^b	14.2 (2.6)	14.3 (2.4)	16.3 (2.1)	13.6 (2.8)	13.8 (2.6)	16.3 (2.1)
Handedness (LQ) ^c	72.8 (27.7)	75.0 (20.2)	75.3 (20.0)	74.5 (28.3)	78.2 (19.4)	74.7 (19.9)
Verbal IQ (WAIS) ^d	102.7 (16.0)	102.9 (14.9)		102.9 (18.0)	98.0 (15.4)	
Onset age (years)	22.4 (6.6)	20.8 (5.1)		22.8 (6.1)	21.1 (6.0)	
Illness duration (years)	7.5 (7.6)	7.8 (9.0)		7.9 (9.0)	7.7 (8.4)	
Total BPRS ^e	45.7 (15.3)	31.1 (10.3)		47.7 (15.5)	32.7 (9.8)	
PANSS general ^e	37.6 (13.4)	27.1 (8.7)		39.4 (13.7)	28.4 (7.8)	
PANSS positive ^e	19.4 (7.3)	12.4 (5.3)		20.4 (7.2)	12.8 (5.4)	
PANSS negative ^e	17.1 (6.6)	12.5 (5.4)		18.2 (7.4)	14.1 (5.7)	
Schizophrenia, paranoid	6	12		8	14	
Schizophrenia, undifferentiated	4	10		6	16	
Schizophrenia, catatonic					1	
Schizophrenia, residual					1	
Schizoaffective, depressed	3	5		2	7	
Schizoaffective, bipolar	5	2		4	4	
Psychosis NOS	1	3		1	4	
Unmedicated (>14 days)	9	11		11	18	
Medicated (atypical antipsychotics)	10	21		10	29	

^aAH: auditory hallucinators; NH: nonhallucinators; HC: healthy controls. Group overlap across paradigms: AH, *n* = 14; NH, *n* = 30; HC, *n* = 42. ^bRM, WM: AH, NH differ significantly from HC (all *p* ≤ .002). ^cLQ: Laterality quotient (Oldfield, 1971) can vary between -100.0 (completely left-handed) and +100.0 (completely right-handed). ^dRM: AH, *n* = 11; NH, *n* = 22; WM: AH, *n* = 14; NH, *n* = 29. ^eRM: AH differ significantly from NH (total BPRS, PANSS general, PANSS positive, all *p* ≤ .001; PANSS negative, *p* = .01); WM: AH (*n* = 18) differ significantly from NH (*n* = 45; total BPRS, PANSS general, PANSS positive, all *p* ≤ .001; PANSS negative, *p* = .03).

All patients met DSM-IV (American Psychiatric Association, 1994) criteria for schizophrenia (paranoid, *n* = 26; undifferentiated, *n* = 22; catatonic, *n* = 1; residual, *n* = 1), schizoaffective disorder (depressed type, *n* = 11; bipolar type, *n* = 9), schizophreniform (*n* = 1), or psychosis not otherwise specified (*n* = 5; the clinical profile of these patients consisted of core psychotic symptoms, including thought disorder, disorganized thinking, paranoid thoughts, and hallucinations, placing them clearly in the range of schizophrenia spectrum disorders). Diagnoses were determined by consensus among senior diagnosticians, clinicians, and clinical research interviewers (Master's level or above) on the basis of a semistructured interview (Nurnberger et al., 1994), which included items from commonly used instruments (SAPS, Scale for the Assessment of Negative Symptoms [SANS], Structured Clinical Interview for DSM-III-R: Patient Edition [SCID-P]; Andreasen, 1983, 1984; Spitzer, Williams, Gibbon, & First, 1990). Thirty-two patients (43%) did not receive antipsychotic medications for at least 14 days before testing. The remaining 43 patients were treated with risperidone (*n* = 11), aripiprazole (*n* = 10), ziprasidone (*n* = 9), olanzapine (*n* = 6), quetiapine (*n* = 4), or clozapine (*n* = 3), with chlorpromazine equivalents ranging from 67 to 1067 mg/day (Woods, 2003). The number of medicated and unmedicated patients did not differ between AH and NH subgroups for either paradigm, $\chi^2(1) \leq 1.18$, *p* ≥ .28.

Patients were compared to 46 healthy volunteers who were recruited from the New York metropolitan area for a payment of US\$15/h and who were without current or past psychopathology based on a standard screening interview (Structured Clinical Interview for DSM-IV Axis-I Disorders–Non-patient Edition [SCID-NP]; First, Spitzer, Gibbon, & Williams, 1996).

All participants had normal or corrected-to-normal vision. The ethnic composition (63 White, 19 Black, 9 Asian, 1 Native American, 17 more than one race, 12 unknown) was representative for the New York region with similar ratios for each subgroup. The experi-

mental protocol was approved by the institutional review board and was undertaken with the understanding and written consent of each participant.

Stimuli and Procedure

The RM paradigm consisted of continuous recognition memory tasks with common words or unknown faces, as detailed elsewhere (Kayser et al., 2010). Briefly, words were 320 English nouns and faces were 320 black-and-white photographs (160 male) taken from a college yearbook. A series of word (or face) stimuli were foveally presented for 500 ms (constant 2.5-s stimulus onset asynchrony) with 114 stimuli per block and 4 blocks per task (912 trials total). Participants indicated for each item whether it was new (never presented in the series) or old (presented previously) by pressing one of two buttons on a response pad. Participants were instructed to respond to every stimulus as quickly and accurately as possible. There was no overlap between blocks for item repetitions.

Building on our previous implementation of a word serial position task (Kayser et al., 2006), the WM paradigm consisted of two serial position tests (27 trials per block, 4 blocks per task, 216 total trials) with the same word or face stimuli used in the RM paradigm. In each trial, a series of three words (or faces) was presented (500 ms, 1.5-s stimulus onset asynchrony) and, after a 3-s delay, one of the items was repeated (probe). Participants indicated the position of the probe item during the encoding series by pressing one of three buttons after a fixed 1.5-s delay.

Data Acquisition, Recording, and Artifact Procedures

Employing an extended 10-20 system (e.g., Jurcak, Tsuzuki, & Dan, 2007; Pivik et al., 1993) 67-channel montage, continuous

EEG was acquired with a 24-bit Biosemi ActiveTwo system (256 samples/s; DC–128 Hz) using a Lycra stretch electrode cap and external electrodes for nose tip (used as off-line reference) and for monitoring bipolar eye activity (left and right outer canthi; above and below right eye). Volume-conducted blink artifacts were effectively removed from the continuous EEG by means of spatial PCA (NeuroScan, 2003; for further details, see Kayser et al., 2010).

For the RM paradigm, 2-s epochs (300 ms prestimulus baseline) were extracted, low-pass filtered at 20 Hz (–24 dB/octave), corrected for horizontal eye movements, and rigorously screened for residual artifacts (Kayser & Tenke, 2006b) or electrode bridging (Tenke & Kayser, 2001). Artifactual or bridged surface potentials were replaced by spherical spline interpolation (Perrin, Pernier, Bertrand, & Echallier, 1989) using the available artifact-free data if possible (i.e., when less than 25% of all EEG channels contained an artifact); otherwise, a trial was rejected. Stimulus-locked ERPs were computed from correct, artifact-free trials separately for each task (i.e., words or faces) but across all other conditions (i.e., old, new, and filler items; cf. Kayser et al., 2010) to maximize the signal-to-noise ratio for the targeted N1 component. The means for the number of trials used to compute these ERP averages ($M \pm SD$, HC vs. NH vs. AH) were 349 ± 41 vs. 333 ± 41 vs. 317 ± 39 for faces, and 372 ± 73 vs. 350 ± 36 vs. 333 ± 42 for words (range 207–447).

For the WM paradigm, 9-s epochs (1.5 s prestimulus baseline of the first encoding item) were processed in an analogous fashion, although both correct and incorrect trials were included. Shorter 1.5-s subepochs (–102–1398 ms around stimulus onset of each trial item) were extracted from these long, artifact-free ERPs using only the matching stimulus pairs (i.e., probe and corresponding encoding item), which were separately combined for each task. The means for the number of trials underlying these ERPs ($M \pm SD$, HC vs. NH vs. AH) were 149 ± 28 vs. 116 ± 33 vs. 101 ± 39 for faces, and 157 ± 32 vs. 156 ± 28 vs. 148 ± 31 for words (range 62–202).

Finally, all ERPs were low-pass filtered (12.5 Hz, –24 dB/octave) and baseline-corrected (100 ms before stimulus onset).

Current Source Density, N1 Latency Jitter, and Principal Components Analysis

The averaged, nose-referenced ERP waveforms were transformed into reference-free CSD estimates ($\mu\text{V}/\text{cm}^2$ units) using a spherical spline surface Laplacian (Perrin et al., 1989; for details, see Kayser, 2009; Kayser & Tenke, 2006a; Kayser et al., 2007). CSD estimates are based on the second spatial derivative of the recorded surface potentials and represent the magnitude of the radial (transcranial) current flow entering (sinks) and leaving (sources) the scalp, thereby mapping the direction, location, and intensity of current generators that underlie an ERP topography (Mitzdorf, 1985; Nicholson, 1973; Nunez & Srinivasan, 2006). Apart from representing a common bridge between scalp-recorded EEG and intracranial local field potential recordings (see Tenke & Kayser, 2012, for a review), a distinct advantage of the surface Laplacian methodology is that any EEG reference will yield the same, that is, *unique* CSD waveforms with reduced volume-conducted contributions, yielding more focused components with unambiguous polarity and sharper topographies compared to ERPs (Kayser et al., 2009).

The visual N1 sink¹ to words is strongly left-lateralized in healthy adults, with a maximum over the inferior lateral-parietal scalp locations (e.g., sites P7 and P9 of the extended 10–20 system over visual object processing regions along the occipitotemporal ventral stream; Ungerleider & Haxby, 1994; cf. Kayser et al., 2006, 2007, 2009, 2010), and this distinct left-larger-than-right N1 sink asymmetry is also seen for nose-referenced ERPs (Kayser et al., 1999; Kayser, Fong, Tenke, & Bruder, 2003). In contrast, the visual N1 sink to faces is less asymmetric, with a right-parietal maximum (Kayser et al., 2010), and corresponds to the face-sensitive N170 component (e.g., Rossion & Jacques, 2008). These lateralized N1 sinks are likely associated with the structural processing of words (visual word form area; e.g., Cohen et al., 2000) or faces (fusiform face area; e.g., Halgren, Raij, Marinkovic, Jousmaki, & Hari, 2000) involving inferior occipital-temporal regions (e.g., fusiform gyrus).

ERP component latency jitter, either between conditions or between subjects, is a well-known problem for quantifying ERP peaks (e.g., Möcks, 1986). For each paradigm (RM, WM) and task (face, word), individual CSD waveform topographies were temporally adjusted for N1 sink peak latency to create optimized subepochs (–50–400 ms) focused on N1 sink activity. After determining the most negative deflection for pooled CSD waveforms using sites P7 and P9 (word; 91–223-ms interval) or P7, P8, P9, and P10 (face; 79–223 ms), all 67 CSD waveforms were jointly aligned with the mean N1 sink peak latency for each task (face, word) and paradigm (RM, WM). For example, if an individual N1 sink peaked at 141 ms in the RM paradigm using words (i.e., using the average of the corresponding CSD waveforms at P7 and P9), all CSDs for this participant, task and paradigm, were shifted two samples (approximately +8 ms) so that the N1 sink maximum would match the respective mean N1 sink peak latency for words during the RM paradigm (approximately 149 ms). Local N1 sink maxima were unambiguously determined within these intervals in almost all instances (100% for words, 97.2% for faces); if not, the CSD waveforms were not shifted. The N1 sink latency shift did not significantly differ between AH, NH, and HC groups in either task or paradigm (overall range –51 to +31 ms); RM, $F(2,92) = 1.48$, $p > .23$; WM, $F(2,109) < 1.0$, n.s.

Optimized CSD waveforms were submitted to unrestricted, temporal PCA derived from the covariance matrix, followed by Varimax rotation of the covariance loadings (Kayser & Tenke, 2003, 2006a) to determine common sources of variance related to N1 sink activity and to quantify its amplitude. The input matrices consisted of 116 variables (samples between –50 and 400 ms) and 12,730 or 15,008 observations stemming from 95 or 112 participants, two tasks (face, word) and 67 electrode sites. This

1. The chosen nomenclature for ERP (“N1”) and CSD (“N1 sink”) components is intended to reflect the crucial difference between reference-dependent surface potentials (ERPs) and their reference-independent surface Laplacian transformations (CSDs) as well as their association. Because the polarity of surface potentials depends on the choice of the EEG reference, which is arbitrary, a negative ERP deflection observed with any given reference may easily revert to a different polarity, topography, or both (i.e., become a “different ERP component”) after re-referencing the data (i.e., applying a different EEG reference), thereby rendering ERPs nonunique (e.g., Kayser & Tenke, 2010). In contrast, CSDs are unique (i.e., reference-free) and therefore have an unambiguous polarity and topography. Here, the “N1 sink” term associates the radial current flow entering the scalp (i.e., the “sink”) with the label “N1” typically used for a corresponding ERP component (i.e., the negativity peaking at about 150 ms at site P7 when using a nose reference). Thus, the term “N1” is generally used to refer to reference-dependent ERP data, whereas the term “N1 sink” is used to refer to its reference-free CSD counterpart.




Figure 1. Mean (\pm SEM) accuracy during recognition memory (RM) and working memory (WM) tasks using words or faces. Overall, remembering faces was more difficult than remembering words. Patients performed more poorly than healthy controls (HC), but auditory hallucinators (AH) did not differ significantly from nonhallucinators (NH), except for the word WM task. Significant contrasts between groups for each paradigm and each task are indicated as follows: * $p \leq .05$, ** $p \leq .01$, *** $p \leq .001$.

approach provides a concise, efficient simplification of the temporal pattern and spatial distribution of neuronal generators, yielding reference-independent CSD factors with unambiguous component polarity and topography (Kayser & Tenke, 2006a; Tenke & Kayser, 2012).

Statistical Analysis

For each paradigm, N1 sink factor scores were submitted to repeated measures analyses of variance (ANOVAs) with group (AH, NH, HC) as a between-subjects factor and task (word, face), hemisphere (left, right), and site (P7/8, P9/10) as within-subjects factors. The selection of two homologous inferior-parietal scalp locations was based on prior research (Kayser et al., 2007). Inasmuch as these sites jointly represented N1 sink maxima, main or interaction effects involving site were not considered.

For analyses of overall performance accuracy, repeated measures ANOVAs with group and task were computed, using the d' -like sensitivity measure d_L (Snodgrass & Corwin, 1988) for the RM paradigm (e.g., Kayser et al., 1999, 2010) and percentages of correct responses (chance levels linearly scaled to 50%) for the WM paradigm (Kayser et al., 2006).

As there were no specific hypotheses regarding sex differences and given the resulting small cell sizes for AH, gender was excluded from all analyses (all reported effects were preserved in secondary analyses including gender). Group differences in accuracy or N1 sink amplitude were examined using planned contrasts (BMDP-4V; Dixon, 1992). All reported correlations coefficients are linear (Pearson). A conventional significance level ($p < .05$) was applied for all effects. Effect sizes (η_p^2) are reported for group main effects.

Results

Behavioral Data

In both paradigms, remembering faces was more difficult than remembering words: task main effect: RM, $F(1,92) = 33.2$,

$p < .0001$; WM, $F(1,109) = 494.7$, $p < .0001$ (Figure 1). Patients had poorer performance than controls: group main effect: RM, $F(2,92) = 7.56$, $p = .0009$, $\eta_p^2 = .141$; WM, $F(2,109) = 23.5$, $p < .0001$, $\eta_p^2 = .301$; between group contrasts: RM, HC vs. AH, $F(1,92) = 11.5$, $p = .001$, HC vs. NH, $F(1,92) = 8.93$, $p = .004$; WM, HC vs. AH, $F(1,109) = 35.5$, $p < .0001$, HC vs. NH, $F(1,109) = 30.5$, $p < .0001$, but hallucinators did not differ significantly from nonhallucinators: RM, $F(1,92) < 1.0$, n.s.; WM, $F[1,109] = 2.57$, $p > .11$. The difference in accuracy between the patient groups and controls was smaller for words than faces during the WM paradigm, which resulted in a significant Group \times Task interaction, $F(2,109) = 27.0$, $p < .0001$. A ceiling performance of controls may have contributed to this effect, and no such interaction was observed during the RM paradigm. However, between group contrasts computed in repeated measures ANOVAs separately for each task and paradigm revealed poorer performance of hallucinators compared to nonhallucinators for words during the WM paradigm (see significance brackets in Figure 1).

Electrophysiologic Data

Figure 2a compares the grand mean CSD waveforms of patients and controls at selected left-hemispheric inferior lateral-parietal sites where N1 sink was prominent for words (P7, P9), and at homologous right-hemispheric sites for faces (P8, P10) for both RM and WM paradigms. These corrected (i.e., peak aligned), grand mean CSDs were highly comparable to those not adjusted for N1 sink peak latency (supplementary Figure S1 along with their nose-referenced ERP counterparts). A distinct component structure, consisting of P1 source (approximate peak latency 85 ms), N1 sink (145 ms), and P2 source (240 ms for faces), was observed for the three groups in both paradigms. In all conditions, however, auditory hallucinators revealed marked reductions of N1 sink compared to nonhallucinators and controls, who had almost identical N1 sink amplitudes. In contrast, no obvious group differences were seen for P1 source amplitude. Parallel ANOVAs computed for P1 source factors (RM vs. WM, 98 vs.




Figure 2. N1 sink amplitude (adjusted for peak latency) for auditory hallucinators (AH), nonhallucinators (NH), and healthy controls (HC) during the recognition memory (RM) and working memory (WM) paradigms using words or faces. A: Grand mean current source density (CSD) [$\mu\text{V}/\text{cm}^2$] waveforms at selected inferior lateral-parietal sites for words (left hemisphere sites P7, P9) and faces (right hemisphere sites P8, P10). These locations are highlighted in the inset showing the 67-channel EEG montage (top view, nose up). B: Time courses of Varimax-rotated covariance loadings for PCA factors corresponding to N1 sink (RM: 141 ms peak latency; WM: 148 ms) and associated factor score topographies for each task and group. C: Mean (\pm SEM) N1 sink amplitude at inferior lateral-parietal sites (P7/8, P9/10) comparing AH, NH, and HC participants. Significant contrasts between groups for each paradigm, task, and hemisphere (LH: left; RH: right) are indicated as follows: (*) $p \leq .10$, * $p \leq .05$, ** $p \leq .01$.

105 ms peak latency, 6.1% vs. 5.2% explained variance) yielded no significant effects involving group (all $F_s \leq 1.88$, all $p_s \geq .16$).

As intended, the unrestricted PCA for these CSD waveforms yielded optimized factors representing N1 sink activity, with highly comparable time courses of factor loadings (RM vs. WM, peak latencies 141 vs. 148 ms, explained variance 16.4% vs. 14.8%; Figure 2b). The corresponding factor score topographies of N1 sink were, for each paradigm, restricted to posterior, inferior-lateral regions, revealing a robust left-larger-than-right hemisphere asymmetry for words and the opposite, less robust asymmetry for faces (Figure 2b). The overall N1 sink amplitude was substantially reduced for auditory hallucinators, although the task-dependent asymmetry was nevertheless preserved.

These observations were confirmed by the statistical analyses. Significant group main effects (RM, $F[2,92] = 3.13$, $p < .05$, $\eta_p^2 = .064$; WM, $F[2,109] = 3.64$, $p = .03$, $\eta_p^2 = .063$) stemmed from smaller N1 sink amplitudes for auditory hallucinators compared to nonhallucinators and healthy controls, who did not differ from each other (between group contrasts: RM, AH vs. NH, $F[1,92] = 3.90$, $p = .05$, AH vs. HC, $F[1,92] = 6.05$, $p = .02$, NH vs. HC, $F[1,92] < 1.0$, n.s.; WM, AH vs. NH, $F[1,109] = 5.96$, $p = .02$, AH vs. HC, $F[1,109] = 6.30$, $p = .01$, NH vs. HC, $F[1,109] < 1.0$, n.s.). There were also highly significant main effects of task (RM, $F[1,92] = 36.4$, $p < .0001$; WM, $F[1,109] = 49.3$, $p < .0001$) and hemisphere (RM, $F[1,92] = 11.7$, $p < .001$; WM, $F[1,109] = 8.75$, $p = .004$), as well as highly significant Task \times Hemisphere interactions (RM, $F[1,92] = 54.0$, $p < .0001$; WM, $F[1,109] = 59.6$, $p < .0001$), but none of these effects interacted with group (all $F_s \leq 1.88$, all $p_s > .15$). Larger N1 sink amplitudes to words than faces and over left than right hemisphere sites were seen for all groups in both paradigms (Figure 2c). These task and hemisphere effects were moderated by a robust left-greater-than-right asymmetry for words, and a smaller right-greater-than-left asymmetry for faces. Notwithstanding the absence of significant interactions of group with task or hemisphere, contrasts between groups separately computed for each paradigm, task, and hemisphere indicated that differences in N1 sink amplitude between auditory hallucinators and nonhallucinators, as well as auditory hallucinators and healthy controls, were most robust over the left hemisphere for words.

To address the potential confound of differences in severity between hallucinators and nonhallucinators (cf. Table 1, BPRS and PANSS), analogous repeated measures analyses of covariance were computed for patients with PANSS ratings, using the total BPRS score as a between-subjects covariate. All critical effects reported above were maintained in these analyses, which is not surprising given that the total BPRS score and all PANSS subscales did not substantially covary with N1 sink amplitude (correlation range: RM, $n = 51$, $-.08 \leq r \leq .21$, all $p_s \geq .14$; WM, $n = 63$, $-.05 \leq r \leq .20$, all $p_s \geq .20$). Furthermore, additional repeated measures ANOVAs with patients grouped into those with a low (<35) or high (≥ 35) total BPRS score (median split) did not reveal any group differences in N1 sink amplitude: RM, $F(1,49) < 1.0$, n.s.; WM, $F(1,61) < 1.0$, n.s.

PCA-based estimates of N1 sink amplitude were highly comparable across paradigms, as revealed by correlations computed for participants who provided data from both paradigms (AH, $n = 14$, $.81 \leq r \leq .98$; NH, $n = 30$, $.87 \leq r \leq .92$; HC, $n = 42$, $.89 \leq r \leq .92$; all $p_s < .001$). To increase statistical power, data stemming from the two paradigms were therefore combined into a single N1 sink estimate by using either the available N1 sink estimate or the pooled mean across paradigms, exploiting that PCA factor scores represent a standardized measure (the overall

mean across participants, tasks, and sites is 0 with a standard deviation of 1). As expected, the repeated measures ANOVA for the combined sample (AH, $n = 26$; NH, $n = 49$; HC, $n = 46$) confirmed the findings for each paradigm, however, at a more robust significance level: group main effect, $F(2,118) = 4.26$, $p = .02$, $\eta_p^2 = .067$; between group contrasts: AH vs. NH, $F(1,118) = 6.55$, $p = .01$, AH vs. HC, $F(1,118) = 7.44$, $p = .007$, NH vs. HC, $F(1,118) < 1.0$, n.s.

Finally, we addressed the question of whether these findings were specific to auditory hallucinations by also taking into account if patients were experiencing other (i.e., visual, olfactory, somatic/tactile, persecutory) but not auditory hallucinations. Because the small number of patients exhibiting other forms of hallucinations (as indicated by SAPS) prevented forming specific subgroups of comparable size, patients not reporting auditory hallucinations (NH) were regrouped into those reporting any other hallucinations (OH) and those without reporting any hallucinations (WH). The repeated measures ANOVA for the pooled N1 sink estimates of the combined sample (AH, $n = 26$; WH, $n = 23$; OH, $n = 26$; HC, $n = 46$) indicated that auditory hallucinators had significantly reduced N1 sinks (AH: -85 ± 1.22) compared to all other groups (WH: -1.52 ± 1.16 ; OH: -1.66 ± 1.95 ; HC: -1.66 ± 1.40), who did not differ from each other: between group contrasts: AH vs. NH, $F(1,117) = 3.78$, $p = .05$, AH vs. OH, $F(1,117) = 5.90$, $p = .02$, AH vs. HC, $F(1,117) = 7.38$, $p = .008$; all other, $F(1,117) < 1.0$, n.s.

Discussion

When we used visually presented common words or unknown faces, schizophrenia patients showed significantly poorer RM and WM performance than healthy controls, which agrees with evidence of impaired memory processing in schizophrenia (e.g., Barch, 2005; Pelletier, Achim, Montoya, Lal, & Lepage, 2005). Despite poorer performance, the associated sequence of cortical activation was highly comparable in patients and controls, with matching N1 sink peak latencies (approximately 150 ms) and topographies (inferior lateral-parietal maximum), replicating previous findings (Kayser et al., 1999, 2006, 2009). Patients and controls revealed distinct, stimulus-dependent N1 sink asymmetries (left larger than right for words and vice versa for faces). However, auditory hallucinators had markedly reduced N1 sink for words and faces compared to healthy controls and nonhallucinators independent of memory paradigm. Avoiding known limitations of reference-dependent ERP measures (Kayser et al., 2006; Kayser & Tenke, 2010; Tenke & Kayser, 2012), the neuronal generators underlying N1 amplitude were concisely summarized by inferior-parietal sinks and quantified by temporal PCA after aligning the individual CSD waveforms for N1 sink peak latency.

This neurophysiological deficit in early visual processing can not be explained by differences in overall symptom severity, which was unrelated to N1 sink amplitude; instead, higher ratings of severity in auditory hallucinators are likely, at least in part, related to the subgroup classification itself. Furthermore, typical concerns of random effects attributable to small, heterogeneous samples are untenable given the large sample and the lack of subgroup differences in demographic (gender, age, education, handedness, verbal IQ) or clinical variables (illness duration and onset).

Auditory hallucinations, considered a hallmark symptom of schizophrenia (Wible et al., 2009), involve activation in auditory association cortex as well as nonsensory cortical and subcortical regions and interfere with cognitive processing (for a review, see

Allen, Laroi, McGuire, & Aleman, 2008). Although the mechanism responsible for cognitive interference remains unknown, the reduction of N1 sink amplitude to words and faces during WM and RM paradigms has implications for understanding its basis.

First, although N1 reduction to tones has been found in psychotic patients during periods of auditory hallucinations compared to symptom-free intervals (Hubl et al., 2007), it is not limited to the auditory modality but occurs for visual processing as well and appears to be largely independent of specific stimulus features (i.e., words or faces). Patients who are prone to auditory hallucinations also showed decreased activation in temporoparietal regions compared to nonhallucinators during a visually presented verbal WM task (Wible et al., 2009) and showed poorer performance than nonhallucinators in both an auditory and visual word serial position test (Bruder et al., 2011). Thus, any interference caused by experiencing auditory hallucinations is not likely due to direct acoustic interference (i.e., hearing voices). However, visually presented stimuli may be rehearsed using the verbal language system, and interference from the “inner voice” could occur in hallucinators (Ford et al., 2001; Wible et al., 2009). Interestingly, restricted perceptual processing may itself cause hallucinatory symptoms, as seen after depriving healthy individuals of visual experience for several days (Merabet et al., 2004).

Second, N1 sink reduction in hallucinators was present despite an adequate, well-above-chance memory performance across tasks and paradigms, which was comparable to nonhallucinators. Notwithstanding a slightly better performance of nonhallucinators than hallucinators for words during the WM paradigm (i.e., the easiest condition with ceiling effects), the comparable overall performance level across tasks and paradigms indicates that the reduced N1 sink was not merely due to poorer attention or distractibility in hallucinators.

Third, the intact P1 source in hallucinators for unambiguous, high-contrast, and meaningful visual stimuli (i.e., words and faces) used in the recognition memory and working memory paradigms, which differ in these regards from more ambiguous stimuli used for visuospatial discrimination tasks (e.g., Butler et al., 2007; Doniger et al., 2002; Foxe et al., 2001), suggests that the N1 sink reduction did not result from a deficit in earlier visual processing. Rather, it involves interference during a subsequent stage of cognitive processing beginning about 150 ms after word or face onset, likely coinciding with stimulus categorization (Rossion & Jacques, 2008) as opposed to low-level perceptual analysis (Butler et al., 2007). This implies that auditory hallucinations, or the impaired functional processes that cause auditory hallucinations, interfere with the integration of incoming perceptual information as reflected by N1, that is, perceptually driven or bottom-up processing involving the ventral occipito-temporal stream is impaired by top-down modulation or cognitive interference (cf. Twomey, Kawabata Duncan, Price, & Devlin, 2011).

Fourth, the fact that visual N1 to words and faces was fully preserved in those schizophrenia patients without auditory hallucinations does not support the bottom-up processing deficit hypothesis in schizophrenia (e.g., Javitt, 2009), at least not without additional qualification. Instead, this finding may indicate that reductions of early perceptual ERP components for any given sample of schizophrenia patients may be primarily due to top-down cognitive interference in patients prone to auditory hallucinations. In this sense, the present findings may offer a cogent explanation for some of the inconsistency in demonstrating a neurophysiological deficit of early visual processing in schizophrenia using electrophysiological methods. It should also be noted that our

experimental procedure and stimuli mainly probed the ventral (parvocellular) visual processing stream or the “what” pathway (Ungerleider & Haxby, 1994), whereas Javitt (2009) reviewed evidence that reflects visual processing deficits in schizophrenia involving the dorsal (magnocellular) processing stream or the “where” pathway. In any case, this hypothesis regarding early processing deficits in schizophrenia and its relation to specific symptom features is in need of further study.

Last, our findings have implications for the hypothesis that auditory hallucinations involve abnormalities of language-related, left-hemisphere function (e.g., Badcock & Hugdahl, 2012). Although N1 sink reduction was greatest for words and over left inferior-parietal sites where N1 sink was largest, the lack of a significant interaction involving group and hemisphere or task or both does not provide statistical support for this hypothesis. Rather, a bilateral inferior-parietal N1 sink reduction in hallucinators was present to both words and faces.

Neurophysiologic abnormalities are more common or more pronounced in the auditory than visual modality (Egan et al., 1994; Ford et al., 1994; Kayser et al., 2009; Pfefferbaum, Ford, White, & Roth, 1989), leading Ford (1999) wondering whether this may be related “to whatever dysfunction leads to more frequent auditory than visual hallucinations” (p. 668). In fact, our sample did not reveal similar N1 sink reductions for patients experiencing other (e.g., visual, olfactory, somatic/tactile) but not auditory hallucinations, which bolsters the notion that impairments in auditory function play a central role in the etiology of schizophrenia. On the other hand, electrophysiological recording conventions, such as using a linked-ears or -mastoids reference that drastically reduces ERP amplitudes at inferior-parietal sites (cf. Kayser et al., 2003, 2009), may have contributed to the notion of modality-specific ERP reductions in schizophrenia. If the underlying generator of the studied ERP phenomena is in close proximity to the EEG reference, as is the case for secondary visual cortex along the ventral processing stream, the surface potentials as measured at nearby sites (e.g., P7/8, P9/10) will be close to zero. In contrast, neuronal generators within auditory cortex will only marginally be affected by this particular EEG reference choice. For these reasons, it may be easier to demonstrate deficits in auditory rather than visual processing in schizophrenia, even if both modalities were equally affected, because existing group or condition differences will be substantially masked for the visual modality by this EEG reference choice. However, this methodological limitation was averted here by employing a reference-free CSD transform.

Among the shortcomings of previous ERP studies examining the neurophysiology of auditory hallucinations are small sample sizes (e.g., $n = 7$ in Hubl et al., 2007). Although using ERP data of 75 right-handed patients allowed studying a reasonably sized subgroup of 26 patients experiencing auditory hallucinations, the sample was heterogenous with regard to diagnostic criteria and medication status. However, it included 32 unmedicated participants, hallucinators and nonhallucinators did not differ in medication status, and previous research has not linked early neurophysiological deficits to antipsychotic medication (cf. Butler et al., 2007; Rosburg et al., 2008).

These findings support the value of a symptom-based approach for dealing with the heterogeneity of schizophrenia and suggest that visual N1 reductions may not represent a biological marker for schizophrenia per se. Although reduced visual N1 may be dependent on the presence of auditory hallucinations during testing (i.e., a marker of current clinical state), it may also represent a stable trait in

patients prone to auditory hallucinations. Additional research is required to address this state versus trait issue and whether the observed N1 sink reductions for auditory hallucinators are specific

to the visual modality or are also present, and perhaps even more robust, for auditory stimuli (Fuxe et al., 2011; Salisbury et al., 2010).

References

- Allen, P., Laroi, F., McGuire, P. K., & Aleman, A. (2008). The hallucinating brain: A review of structural and functional neuroimaging studies of hallucinations. *Neuroscience and Biobehavioral Reviews*, *32*, 175–191. doi:10.1016/j.neubiorev.2007.07.012
- American Psychiatric Association (1994). *Diagnostic and statistical manual of mental disorders* (4th ed). Washington, DC: Author.
- Andreasen, N. C. (1983). *The Scale for the Assessment of Negative Symptoms (SANS)*. Iowa City, IA: The University of Iowa.
- Andreasen, N. C. (1984). *The Scale for the Assessment of Positive Symptoms (SAPS)*. Iowa City, IA: The University of Iowa.
- Badcock, J. C., & Hugdahl, K. (2012). Cognitive mechanisms of auditory verbal hallucinations in psychotic and non-psychotic groups. *Neuroscience and Biobehavioral Reviews*, *36*, 431–438. doi:10.1016/j.neubiorev.2011.07.010
- Barch, D. M. (2005). The cognitive neuroscience of schizophrenia. *Annual Review of Clinical Psychology*, *1*, 321–353. doi:10.1146/annurev.clinpsy.1.102803.143959
- Bruder, G. E., Alschuler, D. M., Kropmann, C. J., Fekri, S., Gil, R. B., Jarskog, L. F., . . . Wexler, B. E. (2011). Heterogeneity of auditory verbal working memory in schizophrenia. *Journal of Abnormal Psychology*, *120*, 88–97. doi:10.1037/a0021661
- Bruder, G., Kayser, J., Tenke, C., Rabinowicz, E., Friedman, M., Amador, X., . . . Gorman, J. (1998). The time course of visuospatial processing deficits in schizophrenia: An event-related brain potential study. *Journal of Abnormal Psychology*, *107*, 399–411. doi:10.1037/0021-843X.107.3.399
- Bruder, G., Rabinowicz, E., Towey, J., Brown, A., Kaufmann, C. A., Amador, X., . . . Gorman, J. M. (1995). Smaller right ear (left hemisphere) advantage for dichotic fused words in patients with schizophrenia. *American Journal of Psychiatry*, *152*, 932–935.
- Butler, P. D., Martinez, A., Fuxe, J. J., Kim, D., Zemon, V., Silipo, G., . . . Javitt, D. C. (2007). Subcortical visual dysfunction in schizophrenia drives secondary cortical impairments. *Brain*, *130*, 417–430. doi:10.1093/brain/awl233
- Cohen, L., Dehaene, S., Naccache, L., Lehericy, S., Dehaene-Lambertz, G., Henaff, M. A., & Michel, F. (2000). The visual word form area: Spatial and temporal characterization of an initial stage of reading in normal subjects and posterior split-brain patients. *Brain*, *123*, 291–307. doi:10.1093/brain/123.2.291
- Dixon, W. J. (Ed.) (1992). *BMDP statistical software manual: To accompany the 7.0 software release*. Berkeley, CA: University of California Press.
- Doniger, G. M., Fuxe, J. J., Murray, M. M., Higgins, B. A., & Javitt, D. C. (2002). Impaired visual object recognition and dorsal/ventral stream interaction in schizophrenia. *Archives of General Psychiatry*, *59*, 1011–1020. doi:10.1001/archpsyc.59.11.1011
- Egan, M. F., Duncan, C. C., Suddath, R. L., Kirsh, D. G., Mirsky, A. F., & Wyatt, R. J. (1994). Event-related potential abnormalities correlate with structural brain alterations and clinical features in patients with chronic schizophrenia. *Schizophrenia Research*, *11*, 259–271. doi:10.1016/0920-9964(94)90020-5
- First, M. B., Spitzer, R. L., Gibbon, M., & Williams, J. B. W. (1996). *Structured Clinical Interview for DSM-IV Axis-I Disorders—Non-patient Edition (SCID-NP)*. New York: Biometrics Research Department, New York State Psychiatric Institute.
- Ford, J. M. (1999). Schizophrenia: The broken P300 and beyond. *Psychophysiology*, *36*, 667–682. doi:10.1111/1469-8986.3660667
- Ford, J. M., Mathalon, D. H., Kalba, S., Whitfield, S., Faustman, W. O., & Roth, W. T. (2001). Cortical responsiveness during inner speech in schizophrenia: An event-related potential study. *American Journal of Psychiatry*, *158*, 1914–1916. doi:10.1176/appi.ajp.158.11.1914
- Ford, J. M., Roach, B. J., Jorgensen, K. W., Turner, J. A., Brown, G. G., Notestine, R., . . . Mathalon, D. H. (2009). Tuning in to the voices: A multisite fMRI study of auditory hallucinations. *Schizophrenia Bulletin*, *35*, 58–66. doi:10.1093/schbul/sbn140
- Ford, J. M., White, P. M., Csernansky, J. G., Faustman, W. O., Roth, W. T., & Pfefferbaum, A. (1994). ERPs in schizophrenia: Effects of antipsychotic medication. *Biological Psychiatry*, *36*, 153–170. doi:10.1016/0006-3223(94)91221-1
- Fuxe, J. J., Doniger, G. M., & Javitt, D. C. (2001). Early visual processing deficits in schizophrenia: Impaired P1 generation revealed by high-density electrical mapping. *NeuroReport*, *12*, 3815–3820. doi:10.1097/00001756-200112040-00043
- Fuxe, J. J., Yeap, S., Snyder, A. C., Kelly, S. P., Thakore, J. H., & Molholm, S. (2011). The N1 auditory evoked potential component as an endophenotype for schizophrenia: High-density electrical mapping in clinically unaffected first-degree relatives, first-episode, and chronic schizophrenia patients. *European Archives of Psychiatry & Clinical Neuroscience*, *261*, 331–339. doi:10.1007/s00406-010-0176-0
- Green, M. F., Hugdahl, K., & Mitchell, S. (1994). Dichotic listening during auditory hallucinations in patients with schizophrenia. *American Journal of Psychiatry*, *151*, 357–362.
- Haenschel, C., Bittner, R. A., Haertling, F., Rotarska-Jagiela, A., Maurer, K., Singer, W., & Linden, D. E. (2007). Contribution of impaired early-stage visual processing to working memory dysfunction in adolescents with schizophrenia: A study with event-related potentials and functional magnetic resonance imaging. *Archives of General Psychiatry*, *64*, 1229–1240. doi:10.1001/archpsyc.64.11.1229
- Halgren, E., Raji, T., Marinkovic, K., Jousmaki, V., & Hari, R. (2000). Cognitive response profile of the human fusiform face area as determined by MEG. *Cerebral Cortex*, *10*, 69–81. doi:10.1093/cercor/10.1.69
- Hubl, D., Koenig, T., Strik, W. K., Garcia, L. M., & Dierks, T. (2007). Competition for neuronal resources: How hallucinations make themselves heard. *British Journal of Psychiatry*, *190*, 57–62. doi:10.1192/bjp.bp.106.022954
- Hugdahl, K., Loberg, E. M., Jorgensen, H. A., Lundervold, A., Lund, A., Green, M. F., & Rund, B. (2008). Left hemisphere lateralisation of auditory hallucinations in schizophrenia: A dichotic listening study. *Cognitive Neuropsychiatry*, *13*, 166–179. doi:10.1080/13546800801906808
- Hugdahl, K., Loberg, E. M., Specht, K., Steen, V. M. van, Wageningen, H., & Jorgensen, H. A. (2007). Auditory hallucinations in schizophrenia: The role of cognitive, brain structural and genetic disturbances in the left temporal lobe. *Frontiers in Human Neuroscience*, *1*, 6. doi:10.3389/neuro.09.006.2007
- Javitt, D. C. (2009). When doors of perception close: Bottom-up models of disrupted cognition in schizophrenia. *Annual Review of Clinical Psychology*, *5*, 249–275. doi:10.1146/annurev.clinpsy.032408.153502
- Javitt, D. C., Shelley, A. M., Silipo, G., & Lieberman, J. A. (2000). Deficits in auditory and visual context-dependent processing in schizophrenia: Defining the pattern. *Archives of General Psychiatry*, *57*, 1131–1137. doi:10.1001/archpsyc.57.12.1131
- Jurcak, V., Tsuzuki, D., & Dan, I. (2007). 10/20, 10/10, and 10/5 systems revisited: Their validity as relative head-surface-based positioning systems. *NeuroImage*, *34*, 1600–1611. doi:10.1016/j.neuroimage.2006.09.024
- Kay, S. R., Opler, L. A., & Fishbein, A. (1992). *Positive and Negative Syndrome Scale (PANSS) rating manual*. Toronto, Canada: Multihealth System Inc.
- Kayser, J. (2009). *Current source density (CSD) interpolation using spherical splines: CSD toolbox (Version 1.1)*. Retrieved from <http://psychophysiology.cpmc.columbia.edu/Software/CSDtoolbox>
- Kayser, J., Bruder, G. E., Friedman, D., Tenke, C. E., Amador, X. F., Clark, S. C., . . . Gorman, J. M. (1999). Brain event-related potentials (ERPs) in schizophrenia during a word recognition memory task. *International Journal of Psychophysiology*, *34*, 249–265. doi:10.1016/S0167-8760(99)00082-3
- Kayser, J., Bruder, G. E., Tenke, C. E., Stuart, B. K., Amador, X. F., & Gorman, J. M. (2001). Event-related brain potentials (ERPs) in schizophrenia for tonal and phonetic oddball tasks. *Biological Psychiatry*, *49*, 832–847. doi:10.1016/S0006-3223(00)01090-8
- Kayser, J., Fong, R., Tenke, C. E., & Bruder, G. E. (2003). Event-related brain potentials during auditory and visual word recognition memory

- tasks. *Cognitive Brain Research*, *16*, 11–25. doi:10.1016/S0926-6410(02)00205-7
- Kayser, J., & Tenke, C. E. (2003). Optimizing PCA methodology for ERP component identification and measurement: Theoretical rationale and empirical evaluation. *Clinical Neurophysiology*, *114*, 2307–2325. doi:10.1016/S1388-2457(03)00241-4
- Kayser, J., & Tenke, C. E. (2006a). Principal components analysis of Laplacian waveforms as a generic method for identifying ERP generator patterns: I. Evaluation with auditory oddball tasks. *Clinical Neurophysiology*, *117*, 348–368. doi:10.1016/j.clinph.2005.08.034
- Kayser, J., & Tenke, C. E. (2006b). Electrical distance as a reference-free measure for identifying artifacts in multichannel electroencephalogram (EEG) recordings. *Psychophysiology*, *43*, S51.
- Kayser, J., & Tenke, C. E. (2010). In search of the Rosetta Stone for scalp EEG: Converging on reference-free techniques. *Clinical Neurophysiology*, *121*, 1973–1975. doi:10.1016/j.clinph.2010.04.030
- Kayser, J., Tenke, C. E., Gates, N. A., & Bruder, G. E. (2007). Reference-independent ERP old/new effects of auditory and visual word recognition memory: Joint extraction of stimulus- and response-locked neuronal generator patterns. *Psychophysiology*, *44*, 949–967. doi:10.1111/j.1469-8986.2007.00562.x
- Kayser, J., Tenke, C. E., Gates, N. A., Kroppmann, C. J., Gil, R. B., & Bruder, G. E. (2006). ERP/CSD indices of impaired verbal working memory subprocesses in schizophrenia. *Psychophysiology*, *43*, 237–252. doi:10.1111/j.1469-8986.2006.00398.x
- Kayser, J., Tenke, C. E., Gil, R. B., & Bruder, G. E. (2009). Stimulus- and response-locked neuronal generator patterns of auditory and visual word recognition memory in schizophrenia. *International Journal of Psychophysiology*, *73*, 186–206. doi:10.1016/j.ijpsycho.2009.02.003
- Kayser, J., Tenke, C. E., Kroppmann, C. J., Fekri, S., Alschuler, D. M., Gates, N. A., . . . Bruder, G. E. (2010). Current source density (CSD) old/new effects during recognition memory for words and faces in schizophrenia and in healthy adults. *International Journal of Psychophysiology*, *75*, 194–210. doi:10.1016/j.ijpsycho.2009.12.001
- Merabet, L. B., Maguire, D., Warde, A., Alterescu, K., Stickgold, R., & Pascual-Leone, A. (2004). Visual hallucinations during prolonged blindfolding in sighted subjects. *Journal of Neuro-Ophthalmology*, *24*, 109–113. doi:10.1097/00041327-200406000-00003
- Mitzdorf, U. (1985). Current source-density method and application in cat cerebral cortex: Investigation of evoked potentials and EEG phenomena. *Physiological Review*, *65*, 37–100.
- Möcks, J. (1986). The influence of latency jitter in principal component analysis of event-related potentials. *Psychophysiology*, *23*, 480–484. doi:10.1111/j.1469-8986.1986.tb00659.x
- Neuhaus, A. H., Karl, C., Hahn, E., Trempler, N. R., Opgen Rhein, C., Urbanek, C., . . . Dettling, M. (2011). Dissection of early bottom-up and top-down deficits during visual attention in schizophrenia. *Clinical Neurophysiology*, *122*, 90–98. doi:10.1016/j.clinph.2010.06.011
- NeuroScan, Inc. (2003). *SCAN 4.3. Vol. II. EDIT 4.3—Offline analysis of acquired data* (Document number 2203, Revision D). El Paso, TX: Compumedics Neuroscan.
- Nicholson, C. (1973). Theoretical analysis of field potentials in anisotropic ensembles of neuronal elements. *IEEE Transactions on Biomedical Engineering*, *20*, 278–288. doi:10.1109/TBME.1973.324192
- Nunez, P. L., & Srinivasan, R. (2006). *Electric fields of the brain: The neurophysics of EEG*. New York: Oxford University Press. doi:10.1093/acprof:oso/9780195050387.001.0001
- Numberger, J. I., Jr., Blehar, M. C., Kaufmann, C. A., York-Cooler, C., Simpson, S. G., Harkavy-Friedman, J., . . . Reich, T. (1994). Diagnostic interview for genetic studies. Rationale, unique features, and training. NIMH Genetics Initiative. *Archives of General Psychiatry*, *51*, 849–859. doi:10.1001/archpsyc.1994.03950110009002
- Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, *9*, 97–113. doi:10.1046/j.1469-7580.2000.19730335.x
- Pascual-Marqui, R. D., Michel, C. M., & Lehmann, D. (1994). Low resolution electromagnetic tomography: A new method for localizing electrical activity in the brain. *International Journal of Psychophysiology*, *18*, 49–65. doi:10.1016/0167-8760(84)90014-X
- Pelletier, M., Achim, A. M., Montoya, A., Lal, S., & Lepage, M. (2005). Cognitive and clinical moderators of recognition memory in schizophrenia: A meta-analysis. *Schizophrenia Research*, *74*, 233–252. doi:10.1016/j.schres.2004.08.017
- Perrin, F., Pernier, J., Bertrand, O., & Echallier, J. F. (1989). Spherical splines for scalp potential and current density mapping [Corrigenda EEG 02274, 1990, *EEG Clin. Neurophysiol.*, *76*, 565]. *Electroencephalography and Clinical Neurophysiology*, *72*, 184–187. doi:10.1016/0013-4694(89)90180-6
- Pfefferbaum, A., Ford, J. M., White, P. M., & Roth, W. T. (1989). P3 in schizophrenia is affected by stimulus modality, response requirements, medication status, and negative symptoms. *Archives of General Psychiatry*, *46*, 1035–1044. doi:10.1001/archpsyc.1989.01810110077011
- Pivik, R. T., Broughton, R. J., Coppola, R., Davidson, R. J., Fox, N., & Nuwer, M. R. (1993). Guidelines for the recording and quantitative analysis of electroencephalographic activity in research contexts. *Psychophysiology*, *30*, 547–558. doi:10.1111/j.1469-8986.1993.tb02081.x
- Price, C. J. (2000). The anatomy of language: Contributions from functional neuroimaging. *Journal of Anatomy*, *197*, 335–359. doi:10.1046/j.1469-7580.2000.19730335.x
- Price, C. J. (2010). The anatomy of language: A review of 100 fMRI studies published in 2009. *Annals of the New York Academy of Sciences*, *1191*, 62–88. doi:10.1111/j.1749-6632.2010.05444.x
- Rosburg, T., Boutros, N. N., & Ford, J. M. (2008). Reduced auditory evoked potential component N100 in schizophrenia: A critical review. *Psychiatry Research*, *161*, 259–274. doi:10.1016/j.psychres.2008.03.017
- Rossion, B., & Jacques, C. (2008). Does physical interstimulus variance account for early electrophysiological face sensitive responses in the human brain? Ten lessons on the N170. *NeuroImage*, *39*, 1959–1979. doi:10.1016/j.neuroimage.2007.10.011
- Salisbury, D. F., Collins, K. C., & McCarley, R. W. (2010). Reductions in the N1 and P2 auditory event-related potentials in first-hospitalized and chronic schizophrenia. *Schizophrenia Bulletin*, *36*, 991–1000. doi:10.1093/schbul/sbp003
- Snodgrass, J. G., & Corwin, J. (1988). Pragmatics of measuring recognition memory: Applications to dementia and amnesia. *Journal of Experimental Psychology: General*, *117*, 34–50. doi:10.1037/0096-3445.117.1.34
- Spitzer, R. L., Williams, J. B. W., Gibbon, M., & First, M. B. (1990). *Structured Clinical Interview for DSM-III-R: Patient edition*. Washington, DC: American Psychiatric Press.
- Tenke, C. E., & Kayser, J. (2001). A convenient method for detecting electrolyte bridges in multichannel electroencephalogram and event-related potential recordings. *Clinical Neurophysiology*, *112*, 545–550. doi:10.1016/S1388-2457(00)00553-8
- Tenke, C. E., & Kayser, J. (2012). *Generator localization by current source density (CSD): implications of volume conduction and field closure at intracranial and scalp resolutions*. *Clinical Neurophysiology*, in press. doi: 10.1016/j.clinph.2012.06.005
- Twomey, T., Kawabata Duncan, K. J., Price, C. J., & Devlin, J. T. (2011). Top-down modulation of ventral occipito-temporal responses during visual word recognition. *NeuroImage*, *55*, 1242–1251. doi:10.1016/j.neuroimage.2011.01.001
- Ungerleider, L. G., & Haxby, J. V. (1994). ‘What’ and ‘where’ in the human brain. *Current Opinion in Neurobiology*, *4*, 157–165. doi:10.1016/0959-4388(94)90066-3
- van Lutterveld, R., Sommer, I. E., & Ford, J. M. (2011). The neurophysiology of auditory hallucinations: A historical and contemporary review. *Frontiers in Psychiatry*, *2*, 28. doi:10.3389/fpsy.2011.00028
- Wible, C. G., Lee, K., Molina, I., Hashimoto, R., Preus, A. P., Roach, B. J., . . . Lauriello, J. (2009). fMRI activity correlated with auditory hallucinations during performance of a working memory task: Data from the FBIRN consortium study. *Schizophrenia Bulletin*, *35*, 47–57. doi:10.1093/schbul/sbn142
- Woods, S. W. (2003). Chlorpromazine equivalent doses for the newer atypical antipsychotics. *Journal of Clinical Psychiatry*, *64*, 663–667. doi:10.4088/JCP.v64n0607
- Yeap, S., Kelly, S. P., Sehatpour, P., Magno, E., Javitt, D. C., Garavan, H., . . . Foxe, J. J. (2006). Early visual sensory deficits as endophenotypes for schizophrenia: High-density electrical mapping in clinically unaffected first-degree relatives. *Archives of General Psychiatry*, *63*, 1180–1188. doi:10.1001/archpsyc.63.11.1180

(RECEIVED April 25, 2012; ACCEPTED May 15, 2012)

Supporting Information

Additional supporting information may be found in the online version of this article:

Figure S1. Uncorrected (i.e., not adjusted for N1 sink peak latency), grand mean (A) nose-referenced surface potential (SP) [μV] and (B) reference-free current source density (CSD) [$\mu\text{V}/\text{cm}^2$] waveforms at selected inferior lateral-parietal sites for words (left hemisphere sites P7, P9) and faces (right hemisphere sites P8, P10) for auditory hallucinators (AH), nonhallucinators

(NH), and healthy controls (HC) during the recognition memory (RM) and working memory (WM) paradigms using words or faces. Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.




Figure S1. Uncorrected (i.e., not adjusted for N1 sink peak latency), grand mean (A) nose-referenced surface potential (SP) [μ V] and (B) reference-free current source density (CSD) [μ V/cm²] waveforms at selected inferior lateral-parietal sites for words (left hemisphere sites P7, P9) and faces (right hemisphere sites P8, P10) for auditory hallucinators (AH), nonhallucinators (NH), and healthy controls (HC) during the recognition memory (RM) and working memory (WM) paradigms using words or faces.