

Accepted Manuscript

Title: Association of posterior EEG alpha with prioritization of religion or spirituality: A replication and extension at 20-year follow-up

Authors: Craig E. Tenke, Jürgen Kayser, Connie Svob, Lisa Miller, Jorge E. Alvarenga, Karen Abraham, Virginia Warner, Priya Wickramaratne, Myrna M. Weissman, Gerard E. Bruder



PII: S0301-0511(17)30005-4
DOI: <http://dx.doi.org/doi:10.1016/j.biopsycho.2017.01.005>
Reference: BIOPSY 7322

To appear in:

Received date: 2-8-2016
Revised date: 10-12-2016
Accepted date: 15-1-2017

Please cite this article as: Tenke, Craig E., Kayser, Jürgen, Svob, Connie, Miller, Lisa, Alvarenga, Jorge E., Abraham, Karen, Warner, Virginia, Wickramaratne, Priya, Weissman, Myrna M., Bruder, Gerard E., Association of posterior EEG alpha with prioritization of religion or spirituality: A replication and extension at 20-year follow-up. *Biological Psychology* <http://dx.doi.org/10.1016/j.biopsycho.2017.01.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Association of posterior EEG alpha with prioritization of religion or spirituality:
a replication and extension at 20-year follow-up**

Craig E. Tenke, Jürgen Kayser, Connie Svob, Lisa Miller, Jorge E. Alvarenga,
Karen Abraham, Virginia Warner, Priya Wickramaratne,
Myrna M. Weissman, Gerard E. Bruder

¹Division of Cognitive Neuroscience., NYS Psychiatric Institute, New York, NY

²Division of Epidemiology, NYS Psychiatric Institute, New York, NY

³Columbia University, College of Physicians and Surgeons, New York, NY

⁴Columbia University, Teachers College, New York, NY

⁵Columbia University, Mailman School of Public Health, New York, NY

Highlights

- Early ratings of religion/spirituality as highly important identified individuals with greater EEG alpha
- EEG alpha was not related to later ratings or changes in religion/spirituality importance
- Religious denomination change was associated with decreased EEG alpha.
- Current generator patterns of posterior resting EEG alpha were concisely quantified by frequency PCA of current source density spectra.

Abstract

A prior report (Tenke et al. 2013 Biol. Psychol. 94:426-432) found that participants who rated religion or spirituality (R/S) highly important had greater posterior alpha after 10 years compared to those who did not. Participants who subsequently lowered their rating also had prominent alpha, while those who increased their rating did not. Here we report EEG findings 20 years after initial assessment. Clinical evaluations and R/S ratings were obtained from 73 (52 new) participants in a longitudinal study of family risk for depression. Frequency PCA of current source density transformed EEG concisely quantified posterior alpha. Those who initially rated R/S as highly important had greater alpha compared to those who did not, even if their R/S rating later increased. Furthermore, changes in religious denomination were associated with decreased alpha. Results suggest the possibility of a critical stage in the ontogenesis of R/S that is linked to posterior resting alpha.

Keywords: EEG alpha; religion and spirituality; current source density (CSD); Principal Components Analysis (PCA); depression risk; development

1. Introduction

Prominent EEG alpha, which is typically obtained from surface potentials measured at scalp, has been associated with better pharmacologic treatment outcomes for clinical depression (Bruder et al., 2008; Tenke et al., 2011; Ulrich et al., 1986). However, systematic comparisons of resting alpha for reference-dependent (nose, average) EEG with reference-free surface Laplacian or current source density (CSD) transformations have demonstrated considerable pitfalls for spectra derived from surface potentials (SP) that are effectively counteracted by CSD (Kayser and Tenke, 2006, 2015a,b; Tenke and Kayser, 2005, 2015). Compared to SP spectra, CSD spectra not only have sharper topographies (Tenke et al., 2015) but also provide the more consistent and appropriate representation of distributed posterior generators in simulation (Tenke & Kayser, 2015), and thereby yield more reliable estimates of alpha activity at scalp. We found that overall posterior CSD alpha during rest is strongly correlated with prestimulus alpha in a novelty oddball task (Tenke et al., 2015), suggesting that posterior alpha may reflect a stable individual trait that cuts across task boundaries or experimental protocols. Even though posterior alpha has been associated with a family history of MDD (Bruder et al., 2005), it remains to be determined to what extent posterior alpha is (1) related to a vulnerability to depression, (2) a predictor of treatment outcome, or (3) an initial vulnerability that is linked to a subsequent propensity toward recovery.

The personal importance of religion or spirituality (R/S) is associated with protection against depression (Kendler et al., 1997; Miller et al., 1997; Smith et al., 2003), and particularly against recurrence of depression (Miller et al., 2012), which is consistent with an acquired resilience following initial vulnerability to MDD. We previously observed that individuals who differed by personal importance of R/S also differed in the prominence of EEG alpha (Tenke et al., 2013) suggesting that

prominent alpha may be a marker for a cluster of related biological, affective and hedonic predispositions. Those who rated R/S as important at an initial test session had greater EEG alpha at a subsequent assessment ten years later than those who did not (Tenke et al., 2013). Our initial considerations led us to hypothesize that EEG alpha might increase in amplitude as a result of R/S importance. In this scenario, alpha enhancement could be a result of meditation or prayer (Aftanas & Golosheykin, 2005; Cahn & Polich, 2006; Chiesa, & Serretti, 2010), stress reduction (Flannelly et al., 2007; Koenig, 2009; Meisenhelder & Marcum, 2009; Stein et al., 2004), or other practices that might accompany R/S. However, our EEG findings were inconsistent with this interpretation, because we observed neither the enhancement of alpha with newly acquired R/S importance, nor its disappearance when the valuation of R/S decreased.

Another conclusion that might be drawn from the data is that EEG alpha could be a marker for the ontogenesis and time course of a disposition toward R/S, rather than either a result or correlate of R/S itself. However, this simpler conclusion is at best preliminary, because the data were limited to changes in R/S Importance assessed at the same time point at which the EEG was measured. It was thereby possible that hypothesized changes in EEG alpha simply required additional time to develop. Conversely, prominent alpha could itself be the trait that protects against MDD and/or is predictive of a good treatment response, and that individuals with this trait disproportionately value R/S highly. Apart from these conceptual considerations, an additional limitation of the original study was a limited spatial sampling of the alpha topography with a low-density EEG montage (i.e., 13 sites), which might have under-represented the complete posterior generator topography of alpha.

The purpose of the present study was to replicate and extend the previous findings on R/S and EEG alpha. After an additional ten year interval, a larger sample of participants were administered two additional R/S assessments and an EEG recording using a higher density montage (i.e., 72 sites)

to better captures the generator topography of EEG alpha. The additional assessments over time allowed us to distinguish between two competing hypotheses: (1) those who were initially identified as having high R/S importance would continue to show high alpha, regardless of intervening R/S reports (i.e. the initial R/S distinguished between individuals differing in the trait of EEG alpha); or (2) individuals who increased, and subsequently maintained, their self-reports of R/S importance would also develop high amplitude alpha (i.e. prominent alpha is associated with persistent R/S importance).

2. Methods

2.1. Participants

This report included seventy-three adult participants (43 female) who had EEG recordings at year 30 (T30; mean age 43.25 yrs \pm 10.2) as part of a multiwave three-generation longitudinal study of individuals at high and low risk for major depression based on family history (Weissman et al, 1997, 2005, 2006, in press). Twenty-one participants were also included in our previous report of the findings at T20 (Tenke et al., 2013), while the remaining 53 only had EEG data at T30. In the original wave of the study (Weissman et al., 1997), probands with moderate to severe major depressive disorder were selected from outpatient clinics for the psychopharmacologic treatment of mood disorders, and nondepressed, demographically-matched control participants were selected from an epidemiologic sample of adults with no psychiatric history from the same community. The sample was recruited from an urban setting (greater New Haven area, Connecticut, US), and consisted of Caucasian and working or middle class individuals. Clinical assessments were conducted by independent interviewers who were blind to the participant's previous clinical history (personal or family).

Assessments were from waves at years 10 (T10; beginning 1992), 20 (T20; beginning 2002), 25 (T25) and 30 (T30). All procedures had been approved by the institutional review boards at Yale University and at Columbia University/New York State Psychiatric Institute. All participants gave written informed consent. As indicated in Table 1, 46 participants who were at high risk due to parental depression were also characterized by significantly greater lifetime rates of MDD compared to those in the low risk group (Fisher exact test, $p = .003$).

(Insert Table 1 about here)

2.2. Religiosity or spirituality

Assessments included participant responses on religious denomination and attendance (“never” to “once a week or more” on a 5-point scale) at T10, T20, T25 and T30. As indicated in Table 2, participants were predominantly Roman Catholic (70%) when first assessed at T10, which dropped to 51% by T30. A total of 44% of the participants reported a change in religious denomination during this time span. R/S importance was also assessed with the question “How important is religion or spirituality to you?” Response option ranged from 1 to 4 (“not important at all,” “slightly important,” “moderately important,” “very important”). This question showed robust correlations with the widely-used Fetzer Institute full-scale measure of personal spirituality (Idler et al., 2003). The terms “religion” and “spirituality” were both included in this question because they are frequently linked together in studies on health (Koenig et al., 2001; Larson and Larson, 2003; Tsuang et al., 2002). Participant responses were dichotomized as “very important” vs. all other responses based on prior findings (Miller et al., 1997, 2012; Tenke et al., 2013), and will hereafter be referred to as “Important” and “Not Important” for sake of brevity.

(Insert Table 2 about here)

Table 3 shows the R/S/ Importance classification of all participants at assessment time T10 and the later assessments, indicating subgroup classifications (*italics*) and corresponding cell sizes for the resulting subgroups and the cross-tabulation marginals (row and column sums). Only 21 participants provided EEG at wave 20 (*ns* in parentheses). Of the 73 in the present sample, 14 were categorized as “Important” at T10. All except 4 of these changed their prioritization at a later assessment (i.e., they “*Migrated Out*” of their prioritization). However, the total number who prioritized R/S as Important at any assessment increased to 37 including 23 participants initially classified as “Not Important” at T10 (i.e., they “*Migrated In*”). The remaining 36 probands were stable in their expression of faith as “Not Important” (i.e., “*Never*”). To be consistent with the classification used in Tenke et al. (2013), comparisons will be made for these four major subgroupings (*Always, Migrated Out, Migrated In, Never*), or after collapsing *Always* and *Migrated Out* probands.

(Insert Table 3 about here)

Table 4 dissects the composition of these subgroupings by familial risk, gender, and lifetime MDD, with bolded labels reflecting the number of probands included for testing the primary hypotheses (i.e., **Important at T10, Migrate In** and **Never**). Women were no more likely than men to rate R/S as “Important” at any assessment (*Always* plus *Migrate Out* plus *Migrate In*; 21 out of 44 vs. 11 out of 29; Fisher exact test, two-sided, *ns*). Likewise, neither risk of MDD due to family history nor lifetime history of MDD distinguished between these three major groupings.

(Insert Table 4 about here)

2.3. EEG methods

Resting EEG was measured while participants sat quietly during four 2-min periods (order of eyes-open and eyes-closed counterbalanced across participants and assessments) after being instructed to avoid blinking and eye and body movements (fixation cross used for eyes-open condition; e.g., cf. Tenke et al., 2011). Continuous data were acquired at 256 samples/s using a 24-bit EEG recording system (BioSemi, Inc., 2001) with a 72-channel 10/10-system scalp montage (Jurcak et al., 2007; Pivik et al., 1993) including the nose. Data were blink-corrected offline using a spatial, singular value decomposition (NeuroScan, 2003), and segmented into 2-s epochs (75% overlap). Epoched data were screened for electrolyte bridges (Alschuler et al., 2014; Tenke and Kayser, 2001), and affected channels interpolated via spherical splines (Perrin et al., 1989). Channels containing artifacts or noise for any given trial were identified using a reference-free approach to identify isolated EEG channels containing amplifier drift, residual eye activity, muscle or movement-related artifacts for any given trial (Kayser and Tenke, 2006b), which were then replaced by spherical spline interpolations (Perrin et al., 1989) from artifact-free channels (i.e., if fewer than 25% of all channels contained artifact). Artifact detection and electrode interpolation was verified interactively. Finally, epochs exceeding $\pm 100 \mu\text{V}$ in any channel (including uncorrected EOG) were rejected to conform to conventional methods.

2.4. Quantification of posterior alpha generators

Artifact-free EEG epochs were transformed to CSD using a spherical spline Laplacian (spline

flexibility constant $m = 4$, regularization constant $\lambda = 10^{-5}$, 50 iterations; Perrin et al., 1989; Kayser and Tenke, 2006, 2015; cf. Kayser, 2009). The DC offset of each epoch was removed, the EEG was tapered over the entire 2-s duration using a Hanning window (Bendat and Piersol, 1971), and padded with zeros (1 s at each end). The CSD power spectra obtained using a subsequent FFT was thereby 0.25 Hz (Neuroscan, 2003), which is consistent with Tenke et al. (2011; 1-s epochs padded to 4 s; 1024 points/epoch) although with less spectral interpolation. CSD amplitude spectra (square root of power spectra; Tenke and Kayser, 2005) were submitted to unrestricted, covariance-based frequency PCA (fPCA) followed by unrestricted Varimax rotation of the covariance loadings (Kayser and Tenke, 2003).

Figure 1 shows the first four CSD-fPCA factors, comprising 85% of the CSD variance, with all remaining factors accounting for less than 4%. Low- and high-frequency alpha factors were unambiguously identifiable by their spectral loadings waveforms and their condition-related topographies (Tenke & Kayser, 2005, 2015; Tenke et al., 2011, 2015). These factors were subsequently pooled over their posterior maxima over each hemisphere (low-frequency alpha, right hemisphere: PO10, PO8, P8; high-frequency, right hemisphere : PO8, PO4, O2, POz, Oz; homologous sites for left hemisphere).

(Insert Figure 1 about here)

2.5. Statistical method

The analysis strategy precisely followed that used in our previous study of EEG and R/S (Tenke et al., 2013). Posterior alpha CSD was compared using a repeated measures ANOVA with alpha *Factor* (high-frequency alpha, low-frequency alpha), *Hemisphere* (left, right) and *Condition*

(eyes-open, eyes-closed) as within-subject factors and *Importance at T10* (Important, Not Important) as the primary grouping factor. Between-subjects variables of *Gender* (male, female), *Parental Depression* (no parental MDD, parental MDD), and *Lifetime Depression* (MDD, No MDD) were also included as control design factors, and *age* at the T30 EEG session as used a covariate. Previous CSD-fPCA findings have not suggested important distinctions between the two posterior alpha factors as related to MDD treatment response (Tenke et al., 2011) or R/S importance (Tenke et al., 2013). Consequently, effects involving *Factor* were not specifically probed (but see Table S1 in Supplementary Material).

Additional analyses followed directly from the findings of Tenke et al. (2013). In particular, *Always Important* and *Migrate Out* were collapsed into a single group, owing to the observation that both groups had greater alpha compared to *Migrate In*, and because of the small number of *Always Important* participants. Likewise, ANOVA models were supplemented by distribution-free statistics (Mann-Whitney U) to compare overall (mean across conditions) and net (difference between conditions) alpha between the same groups. Finally, participants were classified as having high or low overall alpha amplitude based on the median for those who never reported R/S as important (Tenke et al., 2013). The resulting contingency tables were compared using a Fisher exact test.

Inasmuch as religious concordance within families has itself been linked to protection against MDD (Miller et al., 1997; Jacobs et al., 2012), significant findings involving religious importance were followed up by repeated measures ANOVA directly comparing the two *Migrator Groups* (Migrate Out, Migrate In), with *Denomination Change* (changed, not changed), and *Gender* as additional grouping factors, and with *age* and *attendance* at T30 as a covariates.

3. Results

The repeated measures ANOVA model for Importance at T10 (*Always/Migrate Out vs Migrate In/Never*) revealed a Group main effect ($F[1,58] = 5.20$; $p=.026$), owing to significantly greater overall posterior EEG alpha for the *Always/Migrate Out* group. This difference is prominently illustrated by the overall mean factor score topographies shown in Fig. 2, where *Always/Migrate Out* (Important at T10) group has the greatest alpha. The ANOVA for these three groupings supports this observation with a marginally significant *Importance* main effect ($F[2,51]=3.063$; $p=.055$), and a significant *Condition x Parental Depression x Lifetime Depression* interaction ($F[1,51]=6.337$, $p=.015$).¹ The *Condition* main effect was robust, as expected (greater alpha for eyes-closed than eyes-open; $F[1, 51] = 9.889$, $p=.003$; cf. Fig. 1).

(Insert Figure 2 about here)

Fig. 3 shows the scatter plot of overall alpha factor score amplitude for each of the R/S subgroupings. Alpha amplitude was greatest for *Always/Migrate Out* (i.e., those reporting R/S important at T10). Notably, the two *Migrate Out* participants who resumed their high valuation of R/S importance by T30 (green triangles) had no greater overall alpha than those who did not. In contrast, *Migrate In* participants generally showed low overall alpha amplitude, regardless of whether their acquired importance persisted (blue triangles) or was lost by T30 (green inverted triangle). Consequently, the difference between *Always /Migrate Out* and *Migrate In* is robust (Mann-Whitney $U=68$, $N=37$, $p=.003$), as is the contingency table using the *Never* median as a classification threshold (Fisher 2-tailed exact test $p=.006$).

(Insert Figure 3 about here)

Also evident in Fig. 3 is the observation that of those who reported R/S as important at any time, the five individuals with the highest amplitude alpha were the Migrators who retained their religious denomination (filled symbols). The difference in alpha due to denomination change was evaluated by a repeated measures ANOVA with *Denomination Change* (changed, not changed), *Migrator Group* (*Migrate In*, *Migrate Out*) and *Gender* as between-subject factors and *age* and *attendance* as covariates, yielding a significant effect of *Denomination Change* ($F[1, 19] = 5.061, p = .037$). Moreover, a condition-dependent difference between migrator groups was observed (*Condition* \times *Migrator Group* interaction, $F[1,19] = 6.152, p=.023$) in further qualification of a marginally significant overall *Migrator Group* main effect ($F[1,19] = 3.961, p=.061$).

4. Discussion

4.1 EEG Alpha and R/S Importance: Replication and extension

These findings provide additional support for the linkage between the timecourse of personal R/S importance that unfolds over a lifetime and a persistent individual trait of prominent posterior EEG alpha. The present report extends the time scale of this linkage by an additional ten years in a larger, mostly independent sample of participants. Moreover, by virtue of using a substantially denser EEG montage (72 vs. 13 sites), the effects could be unequivocally matched the unique neuronal generator pattern identified with posterior EEG alpha (Tenke et al., 2011, 2015).

These findings closely parallel those reported for the T20 EEG sample (Tenke et al., 2013). Individuals who initially (T10) reported a high personal importance of R/S had significantly greater posterior resting EEG alpha when measured twenty years later (T30). In contrast, those who only

later reported a high personal R/S importance (*Migrate In*) were characterized by low amplitude alpha. Although these differences were quite robust, the *Migrate In* group for the present sample included a number of individuals with high amplitude alpha (8/23 greater than *Never* median), as compared to substantially fewer (2/12) in the previous study (Tenke et al., 2013).

The number of participants included in both the present and the previous study (Tenke et al., 2013; i.e., EEG recorded at *both* T20 and T30) and who reported R/S as Important at T10 was quite small (Table 3, $n = 4$). However, data from a considerably larger sample with EEG at both assessments (T20 and T30 EEG; alpha comparisons based on the smaller montage) revealed a strong test-retest correlation for overall posterior alpha (Tenke et al., submitted). The evidence is therefore quite strong that R/S Importance at T10 uniquely identifies an individual as having prominent overall EEG alpha independent of any subsequent changes in personal evaluation of R/S Importance.

A sizeable subsample of *Migrate In* participants again changed their appraisal of R/S importance by T30 (Fig. 3, those *Migrate In* who migrated out again at T30 [green inverted triangles]). These cases consistently had low amplitude alpha, with only one exception. However, while there were only two *Migrate Out* cases that changed again (*Migrate Out* who migrated in again at T30 [green triangles]), and neither one had exceptionally high amplitude alpha. Collectively, there is little evidence that the persistence or recency of R/S importance has any impact on posterior alpha. However, migrators with the greatest alpha were those with a stable religious denomination (solid symbols). Inasmuch as concordance of denomination across generations is also protective against depression (Miller et al., 1997; Jacobs et al., 2012), a possible association between religious discord and (low) alpha cannot be ruled out. Further inferences cannot be drawn given the limited sample sizes.

4.2 Individual differences in EEG alpha

EEG alpha is characteristically produced in states of relaxed wakefulness with reduced sensory stimulation (Gloor, 1969), which provides the rationale for applying net alpha (eyes closed minus eyes open) as a measure of deactivation (Bruder et al., 1997; Henriques & Davidson, 1990). Not surprisingly, structured meditation practice has been shown to increase EEG alpha (Cahn & Polich, 2006), suggesting that meditation and compatible forms of prayer share a physiological basis with relaxation and disengagement. Conversely, focused attention or stress-related processes are consistent with alpha blocking. This conceptualization implies that a continuum exists for alpha connecting behavioral states with minimal demands, such as the eyes-open/closed resting EEG, to those in more demanding tasks. However, there are substantial individual differences in EEG alpha (e.g., Klimesch, 1999). While some individuals show appreciable alpha during relatively difficult tasks, others may produce minimal or no alpha while relaxing with eyes closed. These limiting conditions (i.e. floor and ceiling effects) offer at least a partial explanation for the observation that prestimulus task-related alpha (auditory novelty oddball) was strongly correlated with overall alpha, but negatively correlated with net alpha (Tenke et al., 2015). This distinction is particularly important because early R/S Importance was an indicator for overall alpha, rather than net alpha, in this and our previous study (Tenke et al., 2013).

4.3 Religious denomination, R/S importance and EEG alpha

The impact of denomination change on posterior resting alpha could have a simple origin. Inasmuch as Migrants view, or have viewed, R/S as very important, a change in denomination has the capacity to index a stressful life event with implications beyond personal R/S. These implications

can include familial relationships, personal identity, and even the disruption of coping strategies previously accessed by personal R/S (Koenig, 2009). In individuals who happen to already have high amplitude resting alpha, there could be a degradation in the rhythm itself, as well as in restorative effects normally derived from relaxed wakefulness, meditation or prayer. It is even possible that the context of the assessments, being part of a longitudinal study within families, could rekindle old conflicts or stress in some individuals.

The twice verified association of early R/S importance with alpha is more complex. Even though denomination change could contribute to the timecourse of alpha, particularly for some migrants, it is not sufficient to explain the unique relevance of the R/S assessment at T10. This problem prompts a fundamental question about R/S importance: What is special about T10? The difference cannot be the age of the participant, because each of the groups contain participants of comparable ages. It also cannot be linked to individual differences, because the two studies were made up of largely independent samples. One possible explanation is that the question, "How important is religion or spirituality to you?," has changed its meaning or selectivity over time.

Local or cultural changes in the practice of, or attitude about, R/S may correspond to the increasing trend in the United States to self-identify as "spiritual but not religious." This trend could itself have been further distorted in the T20 assessment, owing to their administration after the 9/11 terrorist attacks in New York City, the aftermath of which included higher rates of distress reactions (Stein et al., 2004). In one religious sample, most respondents reported feeling a closer connection to God or significant others (Meisenhelder & Marcum, 2009), although those with higher stress or perceived threat engaged in multiple coping mechanisms. Conversely, the impact of political divisiveness, warfare and perceived religious extremism of all forms could result in an "overloaded" question for many individuals. It is likewise unclear how the now ubiquitous engagement with hand-

held technology might affect resting state brain activity and informal R/S practice by competing for the time required for contemplative silence.

4.4 Limitations of R/S importance item

The R/S importance item was chosen over twenty years ago as a brief measure suitable to the time-constraints of the participants in the ongoing longitudinal study of familial risk for MDD, but still likely to distinguish meaningful differences (cf. Miller et al., 1997, 2012). It has also proven to be able to distinguish groups that differ in brain structure (Miller et al., 2013), function (Peterson et al. 2014; Svob et al., 2016), and EEG (Tenke et al., 2013). The present report affirms the original EEG findings using stronger measures, and unambiguously characterizes posterior EEG alpha as the biomarker identified by R/S importance. However, all of these studies are limited by their reliance on this single item, which conflates religion and spirituality.

Pargament (1999) noted that while religion has traditionally provided the essential context for spirituality, a growing number of people identify as “spiritual, but not religious.” The constructs of religion and spirituality have not only been changing over time, both in the psychological literature and the broader culture, but they have become pitted against each other, as the institutional vs. the individual, or even the “bad” vs. the “good.” Of particular importance to investigations of the relationship between R/S and physical or mental health, he observed, “Many definitions of spirituality are in fact definitions of spiritual well-being.” Koenig (2008) pressed this concern even further, noting that some measures confound spirituality and mental health, leading to tautological results that are easily misunderstood by readers and reviewers unfamiliar with the test instruments.

We identified trait-like differences in posterior resting alpha based on the R/S importance measure at a critical period in the ontogenesis of R/S. These differences parallel those anticipated from the presumed protective role of R/S against MDD, raising the possibility that it may in some way

be a proxy for mental health. Although our R/S importance measure does not explicitly confound spirituality and mental health, the way the question is interpreted by study participants will nevertheless be influenced by the ongoing drift in the meaning of these terms. It is hoped that a better understanding of the religious and spiritual characteristics of these participants, such as those provided by a new battery of more refined measures² in our ongoing work, will provide a clearer characterization of the R/S importance groupings and the assortment of these groupings by EEG alpha.

Acknowledgments

Supported by the John Templeton Foundation. The data collection was funded by NIMH grant MH36197, and the CSD methods by MH36295.

Disclosure

Drs. Tenke, Kayser, Warner, Wickramaratne and Bruder, Mr. Alvarenga, and Ms. Abraham have no disclosures to make. Dr. Svob has received funding from the Templeton Foundation, and Dr. Miller has received funding from the Templeton Foundation, the Klingenstein Fund, and the Pritchard Foundation, and received payment from Oxford University Press. Dr. Weissman has received funding from NIMH, the National Institute on Drug Abuse (NIDA), NARSAD, the Sackler Foundation, the Templeton Foundation, and the Interstitial Cystitis Association, and receives royalties from Oxford University Press, Perseus Books, American Psychiatric Association Press, and MultiHealth Systems.

References

- Aftanas, L., & Golosheykin, S. (2005). Impact of regular meditation practice on EEG activity at rest and during evoked negative emotions. *International Journal of Neuroscience*, 115, 893-909.
- Alschuler, D.M., Tenke, C.E., Bruder, G.E., & Kayser, J. (2014). Identifying electrode bridging from electrical distance distributions: a survey of publicly-available EEG data using a new method. *Clinical Neurophysiology*, 125, 484-490.
- Bendat, J.S., & Piersol, A.G. (1971). Random data: analysis and measurement procedures. New York, NY: Wiley-Interscience.
- BioSemi, Inc. (2001). ActiveTwo - Multichannel, DC amplifier, 24-bit resolution, biopotential measurement system with active electrodes (<http://www.biosemi.com>). Amsterdam, NL: Author
- Bruder, G.E., Fong, R., Tenke, C.E., Leite, P., Towey, J.P., Stewart, J.E., McGrath, P.J., & Quitkin, F.M. (1997). Regional brain asymmetries in major depression with or without an anxiety disorder: a quantitative EEG study. *Biological Psychiatry*, 41, 939-948.
- Bruder, G.E., Tenke, C.E., Warner, V., Nomura, Y., Grillon, C., Hille, J., Leite, P., & Weissman, M.M. (2005). Electroencephalographic measures of regional hemispheric activity in offspring at risk for depressive disorders. *Biological Psychiatry*, 57, 328-335.

Bruder, G.E., Sedoruk, J.P., Stewart, J.W., McGrath, P.J., Quitkin, F.M., & Tenke, C.E. (2008). Electroencephalographic alpha measures predict therapeutic response to a selective serotonin reuptake inhibitor antidepressant: pre- and post-treatment findings. *Biological Psychiatry*, *63*, 1171-1177.

Cahn, B.R., & Polich, J. (2006). Meditation States and Traits: EEG, ERP, and Neuroimaging Studies *Psychological Bulletin*, *132*, 180–211.

Chiesa, A., & Serretti, A. (2010). A systematic review of neurobiological and clinical features of mindfulness meditations. *Psychological Medicine*, *40*, 1239-1252.

Flannelly, K.J., Koenig, H.G., Galek, K., & Ellison, C.G. (2007). Beliefs, mental health, and evolutionary threat assessment systems in the brain. *Journal of Nervous and Mental Disorders*, *195*, 996-1003.

Gloor P. (1969). Hans Berger and the discovery of the electroencephalogram. *Electroencephalography and Clinical Neurophysiology, Supplement 28*, 1–36.

Henriques, J.B., & Davidson, R.J. (1990). Regional brain electrical asymmetries discriminate between previously depressed and healthy control subjects. *Journal of Abnormal Psychology*, *99*, 22–31.

Idler, E.L., Musick, M.A., Ellison, C.G., George, L.K., Krause, N., Ory, M.G., Pargament, K.I., Powell, L.H., Underwood, L.G., & Williams, D.R. (2003). Measuring multiple dimensions of religion and

spirituality for health research: conceptual background and findings from the 1998 General Social Survey. *Research on Aging*, 25, 327–365

Jacobs, M., Miller, L., Wickramaratne, P., Gameroff, M., & Weissman, M.M. (2012). Family religion and psychopathology in children of depressed mothers: ten-year follow-up. *Journal of Affective Disorders*, 136, 320-7.

Jurcak, V., Tsuzuki, D. & Dan, I. (2007). 10/20, 10/10, and 10/5 systems revisited: Their validity as relative head-surface-based positioning systems. *NeuroImage*, 34, 1600–1611.

Kayser J. (2009) Current Source Density (CSD) Interpolation using Spherical Splines: CSD Toolbox, Available at: <<http://psychophysiology.cpmc.columbia.edu/Software/CSDtoolbox>>.

Kayser, J., & Tenke, C.E. (2003). Optimizing PCA methodology for ERP component identification and measurement: theoretical rationale and empirical evaluation. *Clinical Neurophysiology*, 114, 2307-2325.

Kayser, J., & Tenke, C.E. (2006a). Principal components analysis of Laplacian waveforms as a generic method for identifying ERP generator patterns: I. Evaluation with auditory oddball tasks. *Clinical Neurophysiology*, 117, 348-368.

Kayser, J., & Tenke, C.E. (2006b). Electrical distance as a reference-free measure for identifying artifacts in multichannel electroencephalogram (EEG) recordings. *Psychophysiology*,

43, S51 (<http://psychophysiology.cpmc.columbia.edu/mmedia/spr2006/ElecDistArti.pdf>).

Kayser, J., & Tenke, C.E. (2015a). Issues and considerations for using the scalp surface Laplacian in EEG/ERP research: a tutorial review. *International Journal of Psychophysiology*, 97, 189-209.

Kayser, J., & Tenke, C.E. (2015b). Hemifield-dependent N1 and event-related theta/delta oscillations: an unbiased comparison of surface Laplacian and common EEG reference choices. *International Journal of Psychophysiology*, 97, 258-270.

Kendler, K.S., Gardner, C.O., & Prescott, C.A. (1997). Religion, psychopathology, and substance use and abuse; a multimeasure, genetic-epidemiologic study. *American Journal of Psychiatry*, 154, 322-329.

Klimesch W (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: a review and analysis. *Brain Research Reviews*, 29, 169-195.

Koenig, H.G. (2008). Concerns about measuring "spirituality" in research. *The Journal of nervous and mental disease*, 196, 349-355.

Koenig, H.G. (2009). Research on Religion, Spirituality, and Mental Health: A Review *Canadian Journal of Psychiatry*, 54, 283-291.

Koenig HG, McCullough M.E., & Larson D.B. (2001). Handbook of Religion and Health. New York,

Oxford University Press.

Kohls, N., Sauer, S., Offenbacher, M., & Giordano, J. (2011). Spirituality: an overlooked predictor of placebo effects? *Philosophical transactions of the Royal Society B*, 366, 1838-1848.

Larson DB, & Larson SS. (2003). Spirituality 's potential relevance to physical and emotional health: a brief review of quantitative research. *Journal of Psychology and Theology*, 31, 37-51

Meisenhelder, J.B., & Marcum, J.P. (2009). Terrorism, Post-traumatic Stress, Coping Strategies, and Spiritual Outcomes. *Journal of Religion and Health*, 48, 46–57

Miller, L., Warner, V., Wickramaratne, P., & Weissman, M. (1997). Religiosity and depression: ten-year follow-up of depressed mothers and offspring. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36, 1416-25.

Miller, L., Wickramaratne, P., Gameroff, M.J., Sage, M., Tenke, C.E., & Weissman, M.M. (2012). Religiosity and major depression in adults at high risk: A ten-year prospective study. *American Journal of Psychiatry*, 169, 89-94

Miller,L., Bansal, R., Wickramaratne, P., Xoa, X., Tenke, C.E., Weissman, M.M., & Peterson, B.S. (2014). Neuroanatomical Correlates of Religiosity and Spirituality In Adults at High and Low Familial Risk for Depression. *JAMA Psychiatry*, 71, 128-35.

NeuroScan, Inc. (2003). SCAN 4.3 - Vol. II. EDIT 4.3 - Offline analysis of acquired data (Document number 2203, Revision D). Compumedics Neuroscan, El Paso, TX.

Pargament, K.I. (1999). The psychology of religion and spirituality? Yes and no. *The International Journal for the Psychology of Religion*, 9, 3-16.

Perrin F, Pernier J, Bertrand O, & Echallier JF. (1989). Spherical splines for scalp potential and current density mapping. *Electroencephalography and clinical Neurophysiology*, 72:184–7 [Corrigenda EEG 02274, *EEG clin Neurophysiol* 1990, 76, 565].

Peterson, B.S., Wang, Z., Horga, G., Warner, V., Rutherford, B., Klahr, K.W., Graniello, B., Wickramaratne, P., Garcia, F., Yu, S., Hao, X., Adams, P.B., Qian, M., Liu, J., Gerber, A. & Weissman, M.M. (2014) Discriminating risk and resilience endophenotypes from lifetime illness effects in familial major depressive disorder. *JAMA Psychiatry*, 71,136-48

Pivik., R.T., Broughton, R.J., Coppola, R., Davidson, R.J., Fox, N., & Nuwer, M.R. (1993). Guidelines for the recording and quantitative analysis of electroencephalographic activity in research contexts. *Psychophysiology*, 30, 547-58.

Schuster, M.A., Stein, B.D., Jaycox, L.H., Collins, R.L., Marshall, G.N., Elliott, M.N., Zhou, A.J., Kanouse, D.E., Morrison, J.L., & Berry, S.H. (2001). A national survey of stress reactions after the September 11, 2001, terrorist attacks, *New England Journal of Medicine*, 345, 1507-1512.

Smith, T.B., McCullough, M.E., & Poll, J. (2003). Religiousness and depression: evidence for a main effect and the moderating influence of stressful life events. *Psychological Bulletin*, 129, 614-36.

Stein, B.D., Elliott, M.N., Jaycox, L.H., Collins, R.L., Berry, S.H., Klein, D.J., & Schuster, M.A. (2004). A national longitudinal study of the psychological consequences of the September 11, 2001 terrorist attacks: reactions, impairment, and help-seeking. *Psychiatry*, 67, 105-117.

Svob, C., Wang, Z., Weissman, M. M., Wickramaratne, P., & Posner, J. (2016). Religious and spiritual importance moderate relation between default mode network connectivity and familial risk for depression. *Neuroscience Letters*, 634, 94-97.

Tenke, C.E., & Kayser, J. (2001). A convenient method for detecting electrolyte bridges in multichannel electroencephalogram and event-related potential recordings. *Clinical Neurophysiology*, 112, 545-550.

Tenke, C.E., & Kayser, J. (2005). Reference-free quantification of EEG spectra: combining current source density (CSD) and frequency principal components analysis (fPCA). *Clinical Neurophysiology*, 116, 2826-2846

Tenke, C.E., & Kayser, J. (2015). Surface Laplacians (SL) and phase properties of EEG rhythms: simulated generators in a volume-conduction model. *International Journal of Psychophysiology*, 97, 285-298.

Tenke, C.E., Kayser, J., Manna, C.B.G., Fekri, S., Kroppmann, C.J., Schaller, J.D., Alschuler, D.M., & Bruder, G.E. (2011). Current source density measures of EEG alpha predict antidepressant treatment response. *Biological Psychiatry*, *70*, 388-394.

Tenke, C.E., Kayser, J., Miller, L., Warner, V., Wickramaratne, P., Weissman, M.M., & Bruder, G.E. (2013). Neuronal generators of posterior EEG alpha reflect individual differences in prioritizing personal spirituality. *Biological Psychology*, *94*, 426-432.

Tenke, C.E., Kayser, J., Abraham, K., Alvarenga, J.E., & Bruder, G.E. (2015). Posterior EEG alpha at rest and during task performance: comparison of current source density and field potential measures. *International Journal of Psychophysiology*, *97*, 299-309.

Tenke, C.E., Kayser, J., Alvarenga, J.E., Abraham, K., Warner, V., Weissman, M.M., & Bruder, G.E. (submitted). Stability of EEG measures over ten years in offspring at risk for depressive disorders.

Weissman, M.M., Warner, V., Wickramaratne, P., Moreau, D., & Olsson, M. (1997). Offspring of depressed parents: Ten years later. *Archives of General Psychiatry*, *54*, 932–940.

Weissman, M.M., Wickramaratne, P., Nomura, Y., Warner, V., Verdeli, H., Pilowsky, D.J., Grillon, C., & Bruder, G. (2005). Families at high and low risk for depression: a 3-generation study. *Archives of General Psychiatry*, *62*, 29-36.

Weissman, M.M., Wickramaratne, P., Nomura, Y., Warner, V., Pilowsky, D., & Verdeli, H. (2006).

Offspring of depressed parents: 20 years later. *American Journal of Psychiatry*, 163, 1001-1008

Weissman, M.M., Wickramaratne, P., Geleroff, M.J., Warner, V., Pilowsky, D., Kohad, R.G., Verdelli, H., Jamie Skipper, J., & Talati, A. (in press). Offspring of Depressed Parents: 30 Years Later. *American Journal of Psychiatry* doi: 10.1176/appi.ajp.2016.15101327


Figure Captions


Fig. 1. Top. The first four factors extracted by unrestricted CSD-fPCA (85% total variance), including factors with low- (blue) and high-frequency (red) alpha loadings peaks (top), each identified by its peak frequency. Factor labels indicate peak frequency and percent variance accounted for by each factor. Bottom. Corresponding factor score topographies and condition dependency for each factor (anterior up; 72 electrode locations identified by circles). Mean amplitudes for both alpha factors were greater for eyes-closed than eyes-open, with distinctive topographies expected for posterior alpha (Tenke and Kayser, 2005; Tenke et al., 2011, 2013). The lowest (1.25 Hz) and highest (24.0 Hz) frequency factors are characteristic of residual eye activity and facial muscle activity (see Tenke and Kayser, 2005).

Fig. 2. Overall mean alpha topographies (mean of low- and high-frequency factor scores across conditions). Participants who rated R/S Important at T10 (*Always/Migrate Out*) had markedly greater posterior alpha than those who did not. Those who only later rated R/S as Important (*Migrate In*) were not significantly different in alpha amplitude than those who *Never* reported R/S as Important.

Fig. 3. Scatterplot of overall posterior CSD alpha (pooled across *Alpha Factor, Condition, Hemisphere and Site*) for individuals in each of the R/S Importance subgroups. The horizontal line reflects the median for those who *Never* reported R/S as Important. Alpha was greater for individuals reporting R/S as "Important at T10" than for those reporting "Not Important at T10." Those who continued to report R/S as Important (*Always*) were indistinguishable from those who *Migrate Out* at one or more later assessments. Although individuals who *Migrate In* after T10 included participants with very low overall alpha, some cases spanned the alpha range of *Always* and *Migrate Out*. Neither the recency

nor the persistence of an Importance rating were associated with prominent overall alpha for the *Migrate In* group. Individuals who *Migrate Out* at T20-T30 are indicated by red downward arrows unless they also reported R/S as Important at T30 (green upward arrows). Likewise, those who *Migrate In* at T20-T30 are indicated by blue upward arrows unless they also reported R/S as Not Important at T30 (green downward arrows). Open symbols: denomination change between T10 and T30; Black symbols: T10 denomination missing.








Table 1 Lifetime MDD by Parental MDD

| | Lifetime MDD | | |
|---------------|--------------|-----|----|
| | No | Yes | |
| No MDD Parent | 18 | 9 | 27 |
| MDD Parent | 13 | 33 | 46 |
| | 31 | 42 | 73 |

Fisher's exact test: p=.003

Table 2. Religious Denomination

| | T10 | T30 |
|----------------------|--------------|--------------|
| Catholic | 47 | 37 |
| Protestant | 10 | 5 |
| Jewish | 4 | 4 |
| Buddhist/Hindu/Islam | -- | 1 |
| Personal Religious | 2 | 13 |
| Agnostic/Atheist | 4 | 8 |
| Other | -- | 5 |
| | <i>N</i> =67 | <i>N</i> =73 |

Table 3 Subgroups for Prioritizing of R/S Importance at Initial (T10) vs. Later Assessment (\geq T20)

| | | Later Assessment (T20, T25, T30) | | |
|-------------------------|---------------|--|--|--|
| | | Important | Not Important | T10 Totals |
| T10 | Important | <i>Always</i> <i>n = 4 (2)</i> | <i>Migrate Out</i> <i>n = 10 (2)</i> | <i>Always/Migrate Out</i> <i>n = 14 (4)</i> |
| | Not Important | <i>Migrate in</i> <i>n = 23 (8*)</i> | <i>Never</i> <i>n = 36 (9)</i> | <i>Never/Migrate In</i> <i>n = 59 (17)</i> |
| Later Assessment Totals | | <i>Always/Migrate In</i> <i>n = 27 (10)</i> | <i>Never/Migrate Out</i> <i>n = 46 (14)</i> | <i>N = 73 (21)</i> |

Migrators are changes from T10;

A subset of 21 participants (ns in parentheses) also participated in Tenke et al. (2013)

* includes 3 identified as 'Never' in Tenke et al. (2013)

Table 4 Characteristics of Importance Subgroups

| | <i>n</i> | Parent MDD | | Gender | | Lifetime MDD | |
|--------------------------|-----------|------------|-----------|-----------|-----------|--------------|-----------|
| | | No | Yes | Female | Male | No | Yes |
| Important at T10 | 14 | 5 | 9 | 8 | 6 | 5 | 9 |
| <i>Always</i> | 4 | 1 | 3 | 2 | 2 | 1 | 3 |
| <i>Migrate Out</i> | 10 | 4 | 6 | 6 | 4 | 4 | 6 |
| Not Important at T10 | 59 | 22 | 37 | 36 | 23 | 26 | 33 |
| <i>Migrate In</i> | 18 | 8 | 10 | 13 | 5 | 8 | 10 |
| <i>Never</i> | 41 | 14 | 27 | 23 | 18 | 18 | 23 |
| | 73 | 27 | 46 | 44 | 29 | 31 | 42 |

Endnotes

¹ ANOVA interactions were observed involving *Factor* (high frequency alpha, low frequency alpha) and *Gender* (male, female). These are presented as supplementary material (Table S1) and will not be further discussed in this report.

² Our ongoing follow-up study employs a multifaceted survey covering various religiosity (i.e., religious identity, religious commitment, religious engagement, personal relationship with God, intrinsic religiosity, religious coping, support derived from a religious community) and spiritual measures (i.e., gratitude, compassion, forgiveness, self-transcendence, ecological awareness, mind-body practices, altruism, love).